

THE DECATEGORIFICATION OF SUTURED FLOER HOMOLOGY

STEFAN FRIEDL, ANDRÁS JUHÁSZ, AND JACOB RASMUSSEN

ABSTRACT. We define a torsion invariant T for every balanced sutured manifold (M, γ) , and show that it agrees with the Euler characteristic of sutured Floer homology SFH . The invariant T is easily computed using Fox calculus. With the help of T , we prove that if (M, γ) is complementary to a Seifert surface of an alternating knot, then $SFH(M, \gamma)$ is either 0 or $\mathbb{Z}$ in every Spin^c structure. T can also be used to show that a sutured manifold is not disk decomposable, and to distinguish between Seifert surfaces.

The support of SFH gives rise to a norm z on $H_2(M, \partial M; \mathbb{R})$. Then T gives a lower bound on the norm z , which in turn is at most the sutured Thurston norm x^s . For closed three-manifolds, it is well known that Floer homology determines the Thurston norm, but we show that $z < x^s$ can happen in general. Finally, we compute T for several wide classes of sutured manifolds.

1. INTRODUCTION

Sutured Floer homology is an invariant of balanced sutured manifolds introduced by the second author [Ju06]. It is an offshoot of the Heegaard Floer homology of Ozsváth and Szabó [OS04a], and contains knot Floer homology [OS04c, Ra03] as a special case. The Euler characteristics of these homologies are torsion invariants of three-manifolds. For example, the Euler characteristic of the Heegaard Floer homology HF^+ is given by Turaev's refined torsion [Tu97, Tu02] and the Euler characteristic of knot Floer homology is given by the Alexander polynomial. In this paper, we investigate the torsion invariant which is the Euler characteristic of sutured Floer homology.

To make a more precise statement, we recall some basic facts about sutured Floer homology. Given a balanced sutured manifold (M, γ) and a relative Spin^c structure $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$, the sutured Floer homology is a finitely generated abelian group $SFH(M, \gamma, \mathfrak{s})$. *A priori*, $SFH(M, \gamma, \mathfrak{s})$ is relatively $\mathbb{Z}/2$ graded. To fix an absolute $\mathbb{Z}/2$ grading, we must specify a homology orientation ω of the pair $(M, R_-(\gamma))$; i.e., an orientation of the vector space $H_*(M, R_-(\gamma); \mathbb{R})$. We denote the resulting invariant by $SFH(M, \gamma, \mathfrak{s}, \omega)$.

Following Turaev, we define a torsion invariant $T_{(M, \gamma)}$ for weakly balanced sutured manifolds, which is essentially the maximal abelian torsion of the pair $(M, R_-(\gamma))$. Our construction is generally very close to Turaev's, but we use handle decompositions

of sutured manifolds in place of triangulations. This makes it easier to define a correspondence between lifts and Spin^c structures. $T_{(M,\gamma)}$ is a function which assigns a number to each $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$. The torsion function is well-defined up to a global factor of ± 1 . Again, to fix the sign, we must specify a homology orientation ω of $(M, R_-(\gamma))$. Then we obtain a function $T_{(M,\gamma,\omega)} : \text{Spin}^c(M, \gamma) \rightarrow \mathbb{Z}$.

Theorem 1. *Let (M, γ) be a balanced sutured manifold. Then for any $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ and homology orientation ω of $(M, R_-(\gamma))$,*

$$T_{(M,\gamma,\omega)}(\mathfrak{s}) = \chi(SFH(M, \gamma, \mathfrak{s}, \omega)).$$

Related invariants have also been studied by Goda and Sakasai [GS08] in the case of homology products, and by Wehrli (in preparation).

It is often convenient to combine the torsion invariants of (M, γ) into a single generating function, which we view as an element of the group ring $\mathbb{Z}[H_1(M)]$. To do so, we fix an affine $H_1(M)$ -isomorphism $\iota : \text{Spin}^c(M, \gamma) \rightarrow H_1(M)$, and write

$$\tau(M, \gamma) = \sum_{\mathfrak{s} \in \text{Spin}^c(M, \gamma)} T_{(M,\gamma)}(\mathfrak{s}) \cdot \iota(\mathfrak{s}).$$

The invariant $\tau(M, \gamma)$ is best thought of as a generalization of the Alexander polynomial to sutured manifolds. Like the classical Alexander polynomial, it is well defined up to multiplication by elements of the form $\pm[h]$ for $h \in H_1(M)$. Many properties of the Alexander polynomial have analogues for $\tau(M, \gamma)$.

For example, if Y is a three-manifold with toroidal boundary, then its Alexander polynomial $\Delta(Y)$ is by definition an invariant of $\pi_1(Y)$. Similarly, for $\tau(M, \gamma)$ the following holds.

Proposition 2. *Let (M, γ) be a balanced sutured manifold such that M is irreducible. Then the invariant $\tau(M, \gamma)$ can be computed from the map $\pi_1(R_-(\gamma)) \rightarrow \pi_1(M)$ using Fox calculus.*

It is worth noting that in order to find Example 8.8, we used a computer program that can calculate $\tau(M, \gamma)$ whenever M is a genus two handlebody.

A well-known theorem of McMullen [Mc02] says that $\Delta(Y)$ gives a lower bound on the Thurston norm of Y . There is a natural extension of the Thurston norm to sutured manifolds due to Scharlemann [Sc89]; we recall its definition in Section 7. For $\alpha \in H_2(M, \partial M; \mathbb{R})$, let $x^s(\alpha)$ denote this sutured Thurston norm.

Proposition 3. *Suppose that (M, γ) is a balanced sutured manifold such that M is irreducible. Let $\mathcal{S} \subset H_1(M)$ be the support of $\tau(M, \gamma)$. Then*

$$\max_{s,t \in \mathcal{S}} (s - t) \cdot \alpha \leq x^s(\alpha).$$

Proposition 3 can be proved analogously to McMullen [Mc02] and Turaev [Tu02]. However, we will not do that here since it is an immediate consequence of Theorem 1 and the following result.

Theorem 4. *Suppose that (M, γ) is a balanced sutured manifold such that M is irreducible. Let $S \subset \text{Spin}^c(M, \gamma)$ be the support of $SFH(M, \gamma)$. Then*

$$\max_{\mathfrak{s}, \mathfrak{t} \in S} \langle \mathfrak{s} - \mathfrak{t}, \alpha \rangle \leq x^s(\alpha).$$

Furthermore, equality holds if ∂M consists only of tori.

In analogy with the situation for closed manifolds, it is tempting to guess that one always has equality in Theorem 4, but using an example of Cantwell and Conlon in [CC06] (cf. Proposition 7.12) we show that this is not the case.

The unit Thurston norm ball of a link complement is always centrally symmetric. We demonstrate in Example 8.5 that S and $\mathcal{S}$ can be centrally asymmetric in general.

One final property of τ is that its “evaluation” under the map $\mathbb{Z}[H_1(M)] \rightarrow \mathbb{Z}[H_1(M, R_-(\gamma))]$ is very simple. More precisely, we show the following.

Proposition 5. *Let $p_* : H_1(M) \rightarrow H_1(M, R_-(\gamma))$ be the natural map. Then*

$$p_*(\tau(M, \gamma)) = \pm I_{H_1(M, R_-(\gamma))},$$

where given a group G we define $I_G \in \mathbb{Z}[G]$ to be

$$I_G = \begin{cases} \sum_{g \in G} g & |G| < \infty, \\ 0 & |G| = \infty. \end{cases}$$

For example, suppose K is a knot in a homology sphere, and let (M, γ) be the sutured manifold whose total space is the complement of K and whose boundary contains two sutures parallel to the meridian of K . Then it can be shown that $\tau(M, \gamma) = \Delta_K(t)$. On the other hand, $H_1(M, R_-(\gamma)) = 0$, so $I_{H_1(M, R_-(\gamma))} = 1$. Thus in this case the proposition reduces to the fact that $\Delta_K(1) = \pm 1$.

Definition 1.1. A balanced sutured manifold (M, γ) is a *sutured L -space* if the group $SFH(M, \gamma)$ is torsion-free and is supported in a single $\mathbb{Z}/2$ homological grading.

Examples of such manifolds are easy to find; e.g., if $R \subset S^3$ is a Seifert surface of an alternating link, then we will show in Corollary 6.10 that the sutured manifold complementary to R is a sutured L -space. The next result follows from Proposition 5.

Corollary 6. *If (M, γ) is a sutured L -space, then for each $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ the group $SFH(M, \gamma, \mathfrak{s})$ is either trivial or isomorphic to $\mathbb{Z}$.*

In the last section, we compute the torsion for a variety of examples, including pretzel surface complements, and for all sutured manifolds complementary to Seifert surfaces of knots with ≤ 9 crossings. In all these examples, the sutured Floer homology is easily determined from the torsion. As an application, we give a simple example of a phenomenon first demonstrated by Goda [Go94]: There exist sutured manifolds whose total space is a handlebody, but which are not disk decomposable.

The paper is organized as follows. In section 2, we recall the relevant facts about sutured Floer homology. We furthermore show how sutured Floer homology behaves

under orientation reversal. Section 3 contains the definition of the torsion, and section 5 explains how to compute it using Fox calculus. Section 4 contains the proof of Theorem 1. In section 6, we discuss some algebraic properties of the torsion, including Proposition 5. Section 7 discusses the relation between SFH and the sutured Thurston norm. Finally, section 8 is devoted to examples.

The authors would like to thank Marc Lackenby for pointing out the connection to reference [GS08]. The second author would also like to thank IHÉS for its hospitality during the course of this work, and the Herchel Smith Fund for their generous support.

Conventions. All 3-manifolds are understood to be oriented and compact. All homology groups are with integral coefficients unless otherwise specified. If X is a submanifold of Y , then $N(X)$ denotes an open tubular neighborhood of X in Y .

2. SUTURED FLOER HOMOLOGY

In this section we recall some relevant facts about sutured manifolds and sutured Floer homology. For full details, we refer the reader to [Ju06].

2.1. Balanced sutured manifolds. For our purposes, a *sutured manifold* (M, γ) is a compact oriented 3-manifold M with boundary together with a set $s(\gamma)$ of oriented simple closed curves on ∂M , called *sutures*. We fix a closed tubular neighborhood $\gamma \subset \partial M$ of the sutures, hence γ is a union of pairwise disjoint annuli. Finally, we require that each component of $R(\gamma) = \partial M \setminus \text{Int}(\gamma)$ be oriented, and that this orientation is coherent with respect to $s(\gamma)$. I.e., if δ is a component of $\partial R(\gamma)$ and is given the boundary orientation, then δ must represent the same homology class in $H_1(\gamma)$ as some suture. Define $R_+(\gamma)$ to be the union of those components of $R(\gamma)$ whose orientation is consistent with the orientation on ∂M induced by M , and let $R_-(\gamma) = R(\gamma) \setminus R_+(\gamma)$.

The notion of a sutured manifold is due to Gabai [Ga83]. The description given above is slightly less general than Gabai's, in that we have omitted the possibility of toroidal sutures.

Example 2.1. Let R be a compact oriented surface with no closed components. Then there is an induced orientation on ∂R . Let $M = R \times [-1, 1]$, define $\gamma = \partial R \times [-1, 1]$, finally put $s(\gamma) = \partial R \times \{0\}$. Such a pair (M, γ) is called a *product sutured manifold*.

Example 2.2. Suppose Y is a closed oriented three-manifold. Let $M = Y \setminus \text{Int}(B^3)$, and let $s(\gamma)$ be a neighborhood of the equator in B^3 . We denote the resulting sutured manifold by $Y(1)$.

Example 2.3. Suppose that L is a link in the oriented three-manifold Y . Then the sutured manifold $Y(L) = (M, \gamma)$ is given by $M = Y \setminus N(L)$, and for each component L_0 of L we take $s(\gamma) \cap \partial N(L_0)$ to be two oppositely oriented meridians of L_0 .

Example 2.4. Let L be a null-homologous link in a closed oriented three-manifold Y , and let R be a Seifert surface for L . If $U \simeq \text{Int}(R) \times (-1, 1)$ is a regular neighborhood

of $\text{Int}(R)$, then the complement $M = Y \setminus U$ is a sutured manifold with $\gamma = \partial R \times [-1, 1]$. The curve $s(\gamma)$ is $\partial R \times \{1/2\}$. Then (M, γ) is called the *sutured manifold complementary to R* , and is denoted by $Y(R)$.

Definition 2.1. A *weakly balanced sutured manifold* is a sutured manifold (M, γ) such that for each component M_0 of M we have

$$\chi(R_+(\gamma) \cap M_0) = \chi(R_-(\gamma) \cap M_0).$$

A *balanced sutured manifold* is a weakly balanced sutured manifold (M, γ) such that M has no closed components and the map $\pi_0(\gamma) \rightarrow \pi_0(\partial M)$ is surjective. Finally, we say that (M, γ) is *strongly balanced* if it is balanced and for each component V of ∂M we have $\chi(R_+(\gamma) \cap V) = \chi(R_-(\gamma) \cap V)$.

Balanced sutured manifolds were defined in [Ju06] and strongly balanced sutured manifolds in [Ju08]. The examples given above are all strongly balanced. Since $2\chi(M) = \chi(\partial M) = \chi(R_-(\gamma)) + \chi(R_+(\gamma))$, for a weakly balanced sutured manifold

$$\chi(M, R_-(\gamma)) = \chi(M, R_+(\gamma)) = 0.$$

Sutured Floer homology is only defined for balanced sutured manifolds. However, we can define the torsion for any weakly balanced sutured manifold.

2.2. Spin^c -structures on sutured manifolds. Suppose that (M, γ) is a sutured manifold. Let v_0 be a nowhere zero vector field along ∂M that points into M along $R_-(\gamma)$, points out of M along $R_+(\gamma)$, and on γ is given by the gradient of a height function $s(\gamma) \times [-1, 1] \rightarrow [-1, 1]$.

Definition 2.2. Let v and w be nowhere zero vector fields on M that agree with v_0 on ∂M . We say that v and w are *homologous* if in each component M' of M there is an open ball $B \subset \text{Int}(M')$ such that $v|(M' \setminus B)$ is homotopic to $w|(M' \setminus B)$ rel ∂M through nowhere zero vector fields. We define $\text{Spin}^c(M, \gamma)$ to be the set of homology classes of nowhere zero vector fields v on M such that $v|_{\partial M} = v_0$.

A priori, this definition appears to depend on the choice of v_0 . However, the space of such vector fields is contractible, so there is a canonical identification between equivalence classes coming from different choices of v_0 . In the case of a closed, oriented 3-manifold the definition is equivalent to the standard definition given in terms of bundles (cf. [Tu97]).

Lemma 2.3. $\text{Spin}^c(M, \gamma) \neq \emptyset$ if and only if (M, γ) is weakly balanced. Furthermore, there exists a free and transitive action of $H^2(M, \partial M) \cong H_1(M)$ on the set $\text{Spin}^c(M, \gamma)$.

Proof. An analogous argument as in the proof of [Ju08b, Proposition 3.5] implies that $\text{Spin}^c(M, \gamma) \neq \emptyset$ if and only if (M, γ) is weakly balanced. It follows from obstruction theory that $\text{Spin}^c(M, \gamma)$ is an affine space over $H^2(M, \partial M; \mathbb{Z})$, since nowhere zero vector fields can be thought of as sections of the unit sphere bundle STM . If

$\mathfrak{s}_1, \mathfrak{s}_2 \in \text{Spin}^c(M, \gamma)$, then $\mathfrak{s}_1 - \mathfrak{s}_2$ is the first obstruction to homotoping vector fields representing $\mathfrak{s}_1$ and $\mathfrak{s}_2$.

If v is a representative of $\mathfrak{s}$ and the simple closed curve c represents $h \in H_1(M)$, then an explicit representative of $\mathfrak{s} + h$ can be obtained by Reeb turbularization, which is described in [Tu90, p.639]. $\square$

2.3. Sutured Floer homology. We now sketch the construction of $SFH(M, \gamma)$. Our starting point is a Heegaard diagram adapted to the pair (M, γ) .

Definition 2.4. A *balanced sutured Heegaard diagram*, in short a *balanced diagram*, is a triple (Σ, α, β) , where Σ is a compact oriented surface with boundary and $\alpha = \{\alpha_1, \dots, \alpha_d\}$ and $\beta = \{\beta_1, \dots, \beta_d\}$ are two sets of pairwise disjoint simple closed curves in $\text{Int}(\Sigma)$ such that $\pi_0(\partial\Sigma) \rightarrow \pi_0(\Sigma \setminus \bigcup \alpha)$ and $\pi_0(\partial\Sigma) \rightarrow \pi_0(\Sigma \setminus \bigcup \beta)$ are both surjective.

Note that the restrictions on α and β are equivalent to the conditions that Σ has no closed components and that the elements of α and β are both linearly independent in $H_1(\Sigma)$.

Every balanced diagram (Σ, α, β) uniquely defines a sutured manifold (M, γ) using the following construction. Let M be the 3-manifold obtained from $\Sigma \times [-1, 1]$ by attaching 2-handles along the curves $\alpha_i \times \{-1\}$ for $i = 1, \dots, d$ and along $\beta_j \times \{1\}$ for $j = 1, \dots, d$. The sutures are defined by taking $\gamma = \partial\Sigma \times [-1, 1]$ and $s(\gamma) = \partial\Sigma \times \{0\}$.

Equivalently, (M, γ) can be constructed from the product sutured manifold $R_-(\gamma) \times I$ by first adding d one-handles to $R_-(\gamma) \times \{1\}$, and then d two-handles. The Heegaard surface Σ is the upper boundary of the manifold obtained by adding the one-handles. The α curves are the belt circles of the one-handles, and the β curves are the attaching circles of the two-handles.

The following proposition combines [Ju06, Proposition 2.9] and [Ju06, Proposition 2.13].

Proposition 2.5. *The sutured manifold defined by a balanced diagram is balanced, and for every balanced sutured manifold there exists a balanced diagram defining it.*

If (Σ, α, β) is a balanced diagram for (M, γ) , then the α and β curves define the tori $\mathbb{T}_\alpha = \alpha_1 \times \dots \times \alpha_d$ and $\mathbb{T}_\beta = \beta_1 \times \dots \times \beta_d$ in the symmetric product $\text{Sym}^d(\Sigma)$. We can suppose that $\bigcup \alpha$ and $\bigcup \beta$ intersect transversally. Then $SFH(M, \gamma)$ is the homology of a chain complex whose generators are the intersection points of $\mathbb{T}_\alpha$ and $\mathbb{T}_\beta$. More concretely, an element of $\mathbb{T}_\alpha \cap \mathbb{T}_\beta$ is a set $\mathbf{x} = \{x_1, \dots, x_d\}$, where each x_i is in some $\alpha_j \cap \beta_k$, and each α and β curve is represented exactly once among the x_i 's. Still more concretely, for each permutation $\sigma \in S_d$ we define

$$(\mathbb{T}_\alpha \cap \mathbb{T}_\beta)_\sigma = \{(x_1, \dots, x_d) : x_i \in \alpha_i \cap \beta_{\sigma(i)}, i = 1, \dots, d\}.$$

Then

$$\mathbb{T}_\alpha \cap \mathbb{T}_\beta = \bigcup_{\sigma \in S_d} (\mathbb{T}_\alpha \cap \mathbb{T}_\beta)_\sigma.$$

The differential in the chain complex is defined by counting rigid holomorphic disks in $\text{Sym}^d(\Sigma)$. Since we are mostly interested in the Euler characteristic of SFH , we will have little need to understand these disks; in fact, the only place they appear is in the proof of Proposition 2.12. For the full definition of the differential, the interested reader is referred to [Ju06].

2.4. Orientations and Grading. Next, we consider the homological grading on the sutured Floer chain complex. In its simplest form, this grading is a relative $\mathbb{Z}/2$ grading given by the sign of intersection in $\text{Sym}^d(\Sigma)$ — two generators have the same grading if the corresponding intersection points in $\mathbb{T}_\alpha \cap \mathbb{T}_\beta$ have the same sign. To fix the sign of intersection, or equivalently, to turn this relative $\mathbb{Z}/2$ grading into an absolute one, we must orient $\text{Sym}^d(\Sigma)$, and the tori $\mathbb{T}_\alpha$ and $\mathbb{T}_\beta$.

The orientation of Σ is determined by the orientation of M . To be precise, Σ is always oriented as the boundary of the compression body determined by the α curves, which we view as a submanifold of M . Using this orientation of Σ , we get the product orientation on $\text{Sym}^d(\Sigma)$. If Σ is endowed with a complex structure compatible with its orientation, then the complex orientation on $\text{Sym}^d(\Sigma)$ agrees with the product orientation. However, to get a well-defined $\mathbb{Z}/2$ grading on $SFH(M, \gamma)$, we will always consider $\text{Sym}^d(\Sigma)$ with $(-1)^{d(d-1)/2}$ times the product orientation.

Choosing an orientation of $\mathbb{T}_\alpha$ is the same as choosing a generator of $\Lambda^d(A)$, where $A \subset H_1(\Sigma; \mathbb{R})$ is the d -dimensional subspace spanned by the α 's. Similarly, an orientation of $\mathbb{T}_\beta$ is specified by a choice of generator for $\Lambda^d(B)$, where B is the subspace of $H_1(\Sigma; \mathbb{R})$ spanned by the β 's. To fix the sign of intersection, we must orient the tensor product $\Lambda^d(A) \otimes \Lambda^d(B)$. The choice of orientation can be expressed in terms of a homology orientation for the pair $(M, R_-(\gamma))$.

Lemma 2.6. *Specifying an orientation of $\Lambda^d(A) \otimes \Lambda^d(B)$ is equivalent to specifying an orientation of $H_*(M, R_-(\gamma); \mathbb{R})$.*

Proof. By Proposition 2.5, the pair (M, γ) is homotopy equivalent to a CW pair (X, Y) such that $X \setminus Y$ is a union of d one-cells and d two-cells. Orienting $H_*(M, R_-(\gamma); \mathbb{R})$ is equivalent to orienting $C_*(X, Y; \mathbb{R})$. This complex is supported in dimensions 1 and 2, with $C_1(X, Y; \mathbb{R}) \cong A$ and $C_2(X, Y; \mathbb{R}) \cong B$. $\square$

Definition 2.7. Suppose that (Σ, α, β) is a balanced diagram such that both Σ and $\Lambda^d(A) \otimes \Lambda^d(B)$ are oriented. Then for $\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$ let $m(\mathbf{x})$ be the intersection sign of $\mathbb{T}_\alpha$ and $\mathbb{T}_\beta$ at $\mathbf{x}$.

Now assume that each $\alpha \in \alpha$ and each $\beta \in \beta$ is oriented. If $x \in \alpha \cap \beta$, then let $m(x)$ denote the intersection sign of α and β at x .

Lemma 2.8. *Let (Σ, α, β) be a balanced diagram such that $\Lambda^d(A) \otimes \Lambda^d(B)$ is oriented. Suppose that each $\alpha \in \alpha$ and each $\beta \in \beta$ is oriented such that the product orientations on $\mathbb{T}_\alpha$ and $\mathbb{T}_\beta$ are consistent with the orientation on $\Lambda^d(A) \otimes \Lambda^d(B)$. If*

$\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$, and $x_i \in \alpha_i \cap \beta_{\sigma(i)}$ for some $\sigma \in S_d$, then

$$m(\mathbf{x}) = \text{sgn}(\sigma) \cdot \prod_{i=1}^d m(x_i).$$

Proof. For $1 \leq i \leq d$ let $a_i \in T_{x_i}\alpha_i$ and $b_{\sigma(i)} \in T_{x_i}\beta_{\sigma(i)}$ be positive tangent vectors. Furthermore, let s_i be a positive basis of $T_{x_i}\Sigma$. Then $s = (-1)^{d(d-1)/2} \cdot s_1 \times \dots \times s_d$ is a positive basis of $T_{\mathbf{x}}\text{Sym}^d(\Sigma)$. Following the notations of [Tu01], we have

$$\begin{aligned} \prod_{i=1}^d m(x_i) &= \prod_{i=1}^d \text{sgn}[a_i b_{\sigma(i)} / s_i] = (-1)^{d(d-1)/2} \cdot \text{sgn}[a_1 b_{\sigma(1)} \dots a_d b_{\sigma(d)} / s] = \\ &= \text{sgn}[a_1 \dots a_d b_{\sigma(1)} \dots b_{\sigma(d)} / s] = \text{sgn}(\sigma) \text{sgn}[a_1 \dots a_d b_1 \dots b_d / s] = \text{sgn}(\sigma) \cdot m(\mathbf{x}). \end{aligned}$$

□

2.5. Generators and Spin^c structures. An important property of the sutured Floer chain complex is that it decomposes as a direct sum over Spin^c structures. Definition 4.5 of [Ju06] explains how to assign a Spin^c structure $\mathfrak{s}(\mathbf{x})$ to each $\mathbf{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$ such that if the boundary $\partial\mathbf{x} = \sum a_i \mathbf{y}_i$, where each a_i is non-zero, then $\mathfrak{s}(\mathbf{x}) = \mathfrak{s}(\mathbf{y}_i)$ for all i . The exact mechanics of this assignment do not concern us at the moment, but we will need to know how to compute the difference between the Spin^c structures assigned to two generators.

Given $\mathbf{x}, \mathbf{y} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$, pick a path θ along the α 's from $\mathbf{x}$ to $\mathbf{y}$. More precisely, θ is a singular 1-chain supported on the α 's with $\partial\theta = \sum_{i=1}^d y_i - \sum_{i=1}^d x_i$. Similarly, choose a path η from $\mathbf{x}$ to $\mathbf{y}$ along the β 's. The difference $\theta - \eta$ represents an element of $H_1(\Sigma)$. If θ' is a different path from $\mathbf{x}$ to $\mathbf{y}$ along the α 's, then the difference $\theta - \theta'$ is a linear combination of the α 's in $H_1(\Sigma)$. Similarly, if η' is another path from $\mathbf{x}$ to $\mathbf{y}$, then $\eta' - \eta$ is a linear combination of β 's. Thus $\theta - \eta$ represents a well defined element of $H_1(\Sigma)/L$, where L is the subspace spanned by the α 's and β 's. We write

$$\mathbf{x} - \mathbf{y} = [\theta - \eta] \in H_1(\Sigma)/L \cong H_1(M).$$

Lemma 2.9. [Ju06, Lemma 4.7] *We have $\mathfrak{s}(\mathbf{x}) - \mathfrak{s}(\mathbf{y}) = \mathbf{x} - \mathbf{y}$ in $H_1(M)$.*

Let (M, γ) be a balanced sutured manifold, $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ a relative Spin^c structure, and choose a homology orientation ω for $(M, R_-(\gamma))$. If $(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta})$ is a balanced diagram for (M, γ) , then we introduce the notation

$$\chi(CF(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta}, \mathfrak{s}, \omega)) = \sum_{\{\mathbf{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta : \mathfrak{s}(\mathbf{x}) = \mathfrak{s}\}} m(\mathbf{x}).$$

Lemma 2.10. *If $(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta})$ and $(\Sigma', \boldsymbol{\alpha}', \boldsymbol{\beta}')$ both represent (M, γ) , then*

$$\chi(CF(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta}, \mathfrak{s}, \omega)) = \chi(CF(\Sigma', \boldsymbol{\alpha}', \boldsymbol{\beta}', \mathfrak{s}, \omega)).$$

Proof. Recall that (Σ, α, β) and $(\Sigma', \alpha', \beta')$ can be connected by a sequence of isotopies, handleslides, and stabilizations/destabilizations. By [Pe08], both isotopies and handleslides correspond to isotopies of $\mathbb{T}_\alpha$ and $\mathbb{T}_\beta$ inside $\text{Sym}^d(\Sigma)$. If a pairs of intersection points $\mathbf{x}$ and $\mathbf{y}$ of $\mathbb{T}_\alpha$ and $\mathbb{T}_\beta$ appear/disappear during such an isotopy, then $\mathbf{x}$ and $\mathbf{y}$ can be connected by a topological Whitney disk, and hence $\mathfrak{s}(\mathbf{x}) = \mathfrak{s}(\mathbf{y})$. Invariance under stabilizarion/destabilization follows immediately from Lemma 2.8. $\square$

Definition 2.11. Let (M, γ) be a balanced sutured manifold, $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$, and ω a homology orientation of $(M, R_-(\gamma))$. Then define

$$\chi(SFH(M, \gamma, \mathfrak{s}, \omega)) = \chi(CF(\Sigma, \alpha, \beta, \mathfrak{s}, \omega)),$$

where (Σ, α, β) is any balanced diagram representing (M, γ) .

In practice, it is convenient to combine the Euler characteristics corresponding to different Spin^c structures into a single generating function, which we view as an element of the group ring $\mathbb{Z}[H_1(M)]$. For this fix an affine isomorphism $\iota : \text{Spin}^c(M, \gamma) \rightarrow H_1(M)$ and let

$$(1) \quad \chi(SFH(M, \gamma, \omega)) = \sum_{\mathbf{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta} m(\mathbf{x})[\iota(\mathfrak{s}(\mathbf{x}))].$$

Then $\chi(SFH(M, \gamma, \omega))$ is well-defined up to multiplication by an element in $H_1(M)$, viewed as a unit in $\mathbb{Z}[H_1(M)]$.

2.6. Duality. Let (M, γ) be a balanced sutured manifold, and denote by $(M, -\gamma)$ the same manifold, but with the orientation of the suture $s(\gamma)$ reversed. The effect of this is to reverse the roles of $R_+(\gamma)$ and $R_-(\gamma)$, more precisely, $R_\pm(-\gamma) = R_\mp(\gamma)$. The same thing happens if we reverse the orientation of M . In this subsection we show that the groups $SFH(M, \gamma)$ and $SFH(M, -\gamma)$ are isomorphic, and that they are ‘dual’ to $SFH(-M, \gamma)$ and $SFH(-M, -\gamma)$. This essentially follows the same way as for ordinary Heegaard Floer homology, though it has not appeared in print before in the case of sutured Floer homology.

If $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ is represented by a nowhere vanishing vector field v , then $-v$ defines a Spin^c structure on both $(M, -\gamma)$ and $(-M, \gamma)$. In both cases we denote the homology class of $-v$ by $-\mathfrak{s}$.

Proposition 2.12. *Let (M, γ) be a balanced sutured manifold and let $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$. Then*

$$SFH(M, \gamma, \mathfrak{s}) \cong SFH(M, -\gamma, -\mathfrak{s})$$

as relatively graded groups, and hence

$$SFH(-M, -\gamma, \mathfrak{s}) \cong SFH(-M, \gamma, -\mathfrak{s}).$$

Moreover, $SFH(M, \gamma, \mathfrak{s})$ and $SFH(-M, \gamma, -\mathfrak{s})$ are the homologies of dual chain complexes, so by the universal coefficient theorem

$$SFH(-M, \gamma, -\mathfrak{s}) \cong \text{Hom}(SFH(M, \gamma, \mathfrak{s}), \mathbb{Z}) \oplus \text{Ext}(SFH(M, \gamma, \mathfrak{s})[1], \mathbb{Z}).$$

Proof. First let us fix our orientation conventions. If (Σ, α, β) is a sutured diagram defining (M, γ) , then the orientation of M is given on $\Sigma \times [-1, 1]$ by taking the product orientation, where $[-1, 1]$ is oriented from -1 to 1 . This orientation naturally extends to the attached 2-handles. Then $s(\gamma) = \partial\Sigma \times \{0\}$ is oriented as the boundary of $\Sigma \times \{0\}$, outward normal first. With these choices, $R_{\pm}(\gamma)$ is obtained from $\Sigma \times \{\pm 1\}$ by doing surgery along the feet of the 2-handles.

Given (M, γ) , choose an admissible sutured diagram (Σ, α, β) defining it. Then $(-\Sigma, \alpha, \beta)$ defines $(-M, -\gamma)$, because the product orientation on $\Sigma \times [-1, 1]$ is reversed, furthermore now the suture is $-\partial\Sigma \times \{0\}$.

If we flip the roles of α and β we get the sutured diagram (Σ, β, α) . Call the sutured manifold defined by it (N, ν) . Then we can define a homeomorphism $h: N \rightarrow M$ which maps $\Sigma \times [-1, 1] \subset N$ to $\Sigma \times [-1, 1] \subset M$ using the formula $h(s, t) = (s, -t)$ and extends to the 2-handles naturally. This map h is orientation reversing from N to M , but preserves the orientation of the sutures. Thus $(N, \nu) = (-M, \gamma)$.

Combining the observations of the previous two paragraphs we see that $(-\Sigma, \beta, \alpha)$ defines $(M, -\gamma)$. The chain complexes $CF(\Sigma, \alpha, \beta)$, $CF(-\Sigma, \alpha, \beta)$, $CF(\Sigma, \beta, \alpha)$, and $CF(-\Sigma, \beta, \alpha)$ all have the same generators, namely $\mathbb{T}_{\alpha} \cap \mathbb{T}_{\beta}$. Let $\mathbf{x}, \mathbf{y}$ be generators of $CF(\Sigma, \alpha, \beta)$ that are connected by a rigid pseudo-holomorphic Whitney disc $u: \mathbb{D} \rightarrow \text{Sym}^d(\Sigma)$. Then $-u$ is a pseudo-holomorphic disc connecting $\mathbf{x}$ to $\mathbf{y}$ in $CF(-\Sigma, \alpha, \beta)$ and also in $CF(\Sigma, \beta, \alpha)$, whereas in $CF(-\Sigma, \beta, \alpha)$ the intersection points $\mathbf{x}, \mathbf{y}$ are connected by u . Thus

$$CF(\Sigma, \alpha, \beta) = CF(-\Sigma, \beta, \alpha)$$

and

$$CF(-\Sigma, \alpha, \beta) = CF(\Sigma, \beta, \alpha),$$

and the chain complex $CF(\Sigma, \beta, \alpha)$ is dual to $CF(\Sigma, \alpha, \beta)$.

To get the refined statement involving the Spin^c structures, observe that if $\mathbf{x}$ is a generator of $CF(\Sigma, \alpha, \beta)$ and $\mathbf{x}'$ is the corresponding generator of $CF(\Sigma, \beta, \alpha)$, then $h_*(\mathfrak{s}(\mathbf{x}')) = -\mathfrak{s}(\mathbf{x})$. On the other hand, the Spin^c structure assigned to $\mathbf{x}$ in $CF(\Sigma, \alpha, \beta)$ and in $CF(-\Sigma, \alpha, \beta)$ can be represented by the same vector field on M . Finally, in $CF(-\Sigma, \beta, \alpha)$ each generator is assigned the same Spin^c structure as in $CF(\Sigma, \alpha, \beta)$. $\square$

3. THE DEFINITION OF THE TORSION FUNCTION

In this section, we first define the maximal abelian torsion of a pair of finite CW complexes. Then we define the torsion invariant for weakly balanced sutured 3-manifolds. Our approach follows closely the ideas of Turaev exposed in [Tu97, Tu98, Tu01, Tu02].

3.1. Lifts and Euler structures. Let (X, Y) be a pair of finite k -dimensional CW complexes with $Y \subset X$ and X connected. We write $H = H_1(X)$ and view H as a

multiplicative group. Denote by $\mathcal{C}$ the set of cells in $X \setminus Y$. Let $\pi : \hat{X} \rightarrow X$ be the universal abelian cover of X and write $\hat{Y} = \pi^{-1}(Y)$.

Definition 3.1. A *lift* l from (X, Y) to $(\hat{X}, \hat{Y})$ is a choice for every $c \in \mathcal{C}$ of a cell $l(c)$ in $\hat{X}$ lying over c . Note that if l' is any other lift then for every $c \in \mathcal{C}$ there is an element $g(c) \in H$ such that $l'(c) = g(c) \cdot l(c)$. We say that l and l' are equivalent if

$$\prod_{c \in \mathcal{C}} g(c)^{(-1)^{\dim c}} \in H$$

is trivial. We denote the set of equivalence classes of lifts by $\text{Lift}(X, Y)$.

We now define an action of H on $\text{Lift}(X, Y)$. First suppose that $X \neq Y$. Let $h \in H$ and suppose that $\mathfrak{l} \in \text{Lift}(X, Y)$ is represented by a lift l . Fix an arbitrary cell $c_0 \in \mathcal{C}$ and suppose that $\dim c_0 = i$. Then $h \cdot \mathfrak{l}$ is represented by the lift l' such that $l'(c_0) = h^{(-1)^i} \cdot l(c_0)$ and $l'(c) = l(c)$ for $c \in \mathcal{C} \setminus \{c_0\}$. If $X = Y$, then $|\text{Lift}(X, Y)| = 1$. Then the action of H is trivial on $\text{Lift}(X, Y)$.

The above definition is independent of the choice of c_0 . If $X \neq Y$, then H acts freely and transitively on $\text{Lift}(X, Y)$. In particular, given $\mathfrak{l}_1, \mathfrak{l}_2 \in \text{Lift}(X, Y)$ we get a well-defined element $\mathfrak{l}_1 - \mathfrak{l}_2 \in H$.

Definition 3.2. For each cell $c \in \mathcal{C}$ pick a point $p(c)$ in c . An *Euler chain* for (X, Y) is a one-dimensional singular chain θ in X with

$$\partial\theta = \sum_{c \in \mathcal{C}} (-1)^{\dim c} p(c).$$

Given two Euler chains θ, η we define $\theta - \eta \in H$ to be the homology class of the 1-cycle $\theta - \eta$. Two Euler chains θ, η are called equivalent if $\theta - \eta$ is trivial in H . We call an equivalence class of Euler chains an *Euler structure* and denote the set of Euler structures by $\text{Eul}(X, Y)$.

Note that $\text{Eul}(X, Y) \neq \emptyset$ if and only if $\chi(X, Y) = 0$. Furthermore, if $\text{Eul}(X, Y) \neq \emptyset$ then H acts freely and transitively on $\text{Eul}(X, Y)$.

Definition 3.3. Suppose that $\chi(X, Y) = 0$. Then we define a map

$$E : \text{Lift}(X, Y) \rightarrow \text{Eul}(X, Y)$$

as follows. Pick a point $\hat{p} \in \hat{X}$. Suppose that $\mathfrak{l} \in \text{Lift}(X, Y)$ and choose a lift l representing $\mathfrak{l}$. For every $c \in \mathcal{C}$, connect $\hat{p}$ and a point $\hat{p}(c) \in l(c)$ with an oriented path $\hat{\theta}(c)$ such that $\partial\hat{\theta}(c) = (-1)^{\dim c}(\hat{p}(c) - \hat{p})$. If $\hat{\theta} = \sum_{c \in \mathcal{C}} \hat{\theta}(c)$, then $\theta = \pi(\hat{\theta})$ is an Euler chain since $\chi(X, Y) = 0$. The Euler structure $\mathfrak{e}$ represented by θ only depends on $\mathfrak{l}$, so we define $E(\mathfrak{l}) = \mathfrak{e}$.

If $X \neq Y$ then the map E is an H -equivariant bijection. If $X = Y$ then $\text{Eul}(X, Y)$ is canonically isomorphic to H , and the image of the unique element of $\text{Lift}(X, Y)$ under E is $0 \in H$.

3.2. Torsion of CW complexes. We continue with the notation from the previous section. Let $\mathfrak{l} \in \text{Lift}(X, Y)$ be a lift represented by l . Furthermore, let φ be a ring homomorphism $\mathbb{Z}[H] \rightarrow \mathbb{F}$ to a field $\mathbb{F}$. In this section, we recall the definition of the Reidemeister–Turaev torsion $\tau^\varphi(X, Y, \mathfrak{l}) \in \mathbb{F}$.

Consider the chain complex

$$C_*(X, Y; \mathbb{F}) = C_*(\hat{X}, \hat{Y}; \mathbb{Z}) \otimes_{\mathbb{Z}[H]} \mathbb{F}.$$

Here H acts via deck transformations on $\hat{X}$, and hence on $C_*(\hat{X}, \hat{Y}; \mathbb{Z})$, and H acts on $\mathbb{F}$ via φ . If this complex is not acyclic; i.e., if the twisted homology groups $H_*(X, Y; \mathbb{F})$ do not vanish, then we set $\tau^\varphi(X, Y, \mathfrak{l}) = 0 \in \mathbb{F}$. If the complex is acyclic, then we can define the torsion $\tau^\varphi(X, Y, \mathfrak{l}) \in \mathbb{F} \setminus \{0\}$ as follows.

The cells $\{l(c) : c \in \mathcal{C}\}$ define a basis of $C_*(\hat{X}, \hat{Y}; \mathbb{Z})$ as a complex of free $\mathbb{Z}[H]$ -modules, and hence give a basis for $C_*(X, Y; \mathbb{F})$ as a complex of $\mathbb{F}$ -modules. We now define $\tau^\varphi(X, Y, \mathfrak{l}) \in \mathbb{F}$ to be the torsion of the based acyclic complex $C_*(X, Y; \mathbb{F})$. It is straightforward to check that this is independent of the choice of representative lift corresponding to $\mathfrak{l} \in \text{Lift}(X, Y)$. We refer to [Tu01] for an excellent introduction to the torsion of based complexes.

The torsion $\tau^\varphi(X, Y, \mathfrak{l}) \in \mathbb{F}$ is well-defined up to sign; this indeterminacy comes from the fact that we do not have a canonical ordering of the basis. We can eliminate this indeterminacy by equipping (X, Y) with a homology orientation ω ; i.e., an orientation of the vector space $H_*(X, Y; \mathbb{R}) = \bigoplus_{i \geq 0} H_i(X, Y; \mathbb{R})$. Then we can get a well-defined element $\tau^\varphi(X, Y, \mathfrak{l}, \omega) \in \mathbb{F}$. We refer to [Tu02, Section K] for details.

Now suppose that $\chi(X, Y) = 0$. If $X \neq Y$, then for $\mathfrak{e} \in \text{Eul}(X, Y)$ let

$$\tau^\varphi(X, Y, \mathfrak{e}, \omega) = \tau^\varphi(X, Y, E^{-1}(\mathfrak{e}), \omega).$$

If $X = Y$, then recall that $\text{Eul}(X, Y)$ is canonically identified with H . For $h \in H = \text{Eul}(X, Y)$ let $\tau^\varphi(X, Y, h, \omega) = h$ if ω is the positive orientation of $H_*(X, Y; \mathbb{R}) = 0$, and $\tau^\varphi(X, Y, h, \omega) = -h$ otherwise.

3.3. The maximal abelian torsion of a CW complex. We continue with the notation from the previous sections. In particular, let (X, Y) be a pair of finite CW complexes such that X is connected. Furthermore, let $\mathfrak{l} \in \text{Lift}(X, Y)$ and ω a homology orientation.

Again, we write $H = H_1(X)$ and think of H as a multiplicative group. We let $T = \text{Tor}(H)$ be the torsion subgroup. Given a ring R , we denote by $Q(R)$ the ring which is given by inverting all elements of R that are not zero divisors. We write $Q(H) = Q(\mathbb{Z}[H])$.

A character $\chi : T \rightarrow \mathbb{C}^*$ extends to a ring homomorphism $\chi : \mathbb{Q}[T] \rightarrow \mathbb{C}$, its image is a cyclotomic field $\mathbb{F}_\chi$. Two characters χ_1, χ_2 are called equivalent if $\mathbb{F}_{\chi_1} = \mathbb{F}_{\chi_2}$ and if χ_1 is the composition of χ_2 with a Galois automorphism of $\mathbb{F}_{\chi_1}$ over $\mathbb{Q}$. For any complete family of representatives $\chi_1, \dots, \chi_n$ of the set of equivalence classes of

characters the homomorphism

$$(\chi_1, \dots, \chi_n) : \mathbb{Q}[T] \rightarrow \bigoplus_{i=1}^n \mathbb{F}_{\chi_i}$$

is an isomorphism of rings. We will henceforth identify $\mathbb{Q}[T]$ with $\bigoplus_{i=1}^n \mathbb{F}_{\chi_i}$. Note that under this isomorphism $1 \in \mathbb{Q}[T]$ corresponds to $(1, \dots, 1)$.

Now let F be the free abelian group H/T and pick a splitting $H = F \times T$. Then we have the identifications

$$\mathbb{Q}[H] = \bigoplus_{i=1}^n \mathbb{F}_{\chi_i}[F] \text{ and } Q(H) = \bigoplus_{i=1}^n Q(\mathbb{F}_{\chi_i}[F]).$$

We denote by φ_i the ring homomorphism

$$\mathbb{Z}[H] \rightarrow \bigoplus_{i=1}^n \mathbb{F}_{\chi_i}[F] \rightarrow \mathbb{F}_{\chi_i}[F] \rightarrow Q(\mathbb{F}_{\chi_i}[F]).$$

We now let

$$\tau(X, Y, \mathfrak{l}, \omega) = \sum_{i=1}^n \tau^{\varphi_i}(X, Y, \mathfrak{l}, \omega) \in \bigoplus_{i=1}^n Q(\mathbb{F}_{\chi_i}[F]) = Q(H).$$

Note that $\tau(X, Y, \mathfrak{l}, \omega) \in Q(H)$ is independent of the choices we made (cf. [Tu02, Section K]). Also note that

$$(2) \quad \tau(X, Y, h \cdot \mathfrak{l}, \pm \omega) = \pm h \cdot \tau(X, Y, \mathfrak{l}, \omega).$$

In the following, we write $\tau(X, Y, \mathfrak{l})$ for the set of torsions corresponding to all possible orientations. Also, if $\chi(X, Y) = 0$ and $X \neq Y$, then for $\mathfrak{e} \in \text{Eul}(X, Y)$ we define

$$\tau(X, Y, \mathfrak{e}, \omega) = \tau(X, Y, E^{-1}(\mathfrak{e}), \omega).$$

If $X = Y$ and $\mathfrak{e} \in \text{Eul}(X, Y)$ corresponds to $h \in H$ under the canonical isomorphism $\text{Eul}(X, Y) = H$, then we define $\tau(X, Y, \mathfrak{e}, \omega) = \pm h$, depending on whether ω is the positive or negative orientation of the zero space. Finally, if $\chi(X, Y) \neq 0$, then we set $\tau(X, Y, \mathfrak{e}, \omega) = 0$.

In the coming sections we will often make use of the following two lemmas.

Lemma 3.4. *Let (X, Y) be a pair of finite CW-complexes, and let ω be a homology orientation for (X, Y) . Assume that (X', Y') is a CW-pair obtained from (X, Y) by a simple homotopy s . Then there exists an orientation ω' of (X', Y') and an H -equivariant bijection $b_s : \text{Eul}(X, Y) \rightarrow \text{Eul}(X', Y')$ such that for every $\mathfrak{e} \in \text{Eul}(X, Y)$*

$$\tau(X, Y, \mathfrak{e}, \omega) = \tau(X', Y', b_s(\mathfrak{e}), \omega').$$

Proof. Since $H_*(X', Y'; \mathbb{R}) = H_*(X, Y; \mathbb{R})$, the orientation ω of (X, Y) immediately gives rise to an orientation ω' of (X', Y') . It is sufficient to show the result if X' is obtained from X using an elementary expansion. Suppose that we added an i -cell c and an $(i+1)$ -cell d to X to get X' . Choose an Euler chain θ representing $\mathfrak{e}$. Let

$\delta \subset c \cup d$ be a curve such that $\partial\delta = (-1)^i(p - q)$, where p is the center of c and q is the center of d . Then define $b_s(\mathfrak{e})$ to be the equivalence class of the Euler chain $\theta + \delta$. From here a standard argument shows that $\tau(X, Y, \mathfrak{e}, \omega) = \tau(X', Y', b_s(\mathfrak{e}), \omega')$. $\square$

Lemma 3.5. *Suppose that X is a 2-complex and that $Y \neq \emptyset$. Then for any $\mathfrak{l} \in \text{Lift}(X, Y)$ we have $\tau(X, Y, \mathfrak{l}) \in \mathbb{Z}[H]$.*

Proof. Note that since X is connected and $Y \neq \emptyset$, after collapsing one cells and using Lemma 3.4, we can assume without loss of generality that $X \setminus Y$ contains no 0-cells. Our assumptions then imply that $C_i(X, Y; \mathbb{Z}[H]) = 0$ for any $i \neq 1, 2$. If $\chi(X, Y) \neq 0$, then $H_*(X, Y; \mathbb{F})$ is nontrivial for every $\mathbb{F}$, and the torsion is 0 by definition. If $\chi(X, Y) = 0$, then let A be the matrix representing the boundary map

$$C_2(X, Y; \mathbb{Z}[H]) \rightarrow C_1(X, Y; \mathbb{Z}[H])$$

with respect to a $\mathbb{Z}[H]$ -basis corresponding to $\mathfrak{l} \in \text{Lift}(X, Y)$. Then for any ring homomorphism $\varphi : \mathbb{Z}[H] \rightarrow \mathbb{F}$ we have $\tau^\varphi(M, \gamma, \mathfrak{l}) = \det(\varphi(A)) = \varphi(\det(A))$. It follows immediately from the definitions that $\tau(X, Y, \mathfrak{l}) = \det(A) \in \mathbb{Z}[H]$. $\square$

3.4. Torsion for sutured manifolds. Throughout this section, (M, γ) will be a connected weakly balanced sutured manifold.

Definition 3.6. A *sutured handle complex* $\mathcal{A}$ is a pair of spaces $(A, S \times I)$, where S is a compact surface with boundary, together with a decomposition of $A \setminus (S \times I)$ into 3-dimensional handles $e_1 \cup \dots \cup e_n$, attached to $S \times I$ one after another. More precisely, for $0 \leq i \leq n$ we write $A_i = (S \times I) \cup (e_1 \cup \dots \cup e_i)$ and $S_i = \partial A_i \setminus (S \times \{0\} \cup \partial S \times I)$. Then e_i is smoothly attached to A_{i-1} along $\text{Int}(S_{i-1})$ for $1 \leq i \leq n$.

Let $I(r)$ denote the index of the handle e_r . We say that $\mathcal{A}$ is *nice* if I is non-decreasing and $e_i \cap e_j = \emptyset$ whenever $I(e_i) = I(e_j)$ and $i \neq j$.

Definition 3.7. A *handle decomposition* Z of (M, γ) consists of a sutured handle complex $\mathcal{A}$ and a diffeomorphism $d : A \rightarrow M$ such that $d(S \times \{0\}) = R_-(\gamma)$ and $d(\partial S \times I) = \gamma$. We say that Z is *nice* if $\mathcal{A}$ is nice.

It is straightforward to define $\text{Lift}(Z)$, $\text{Eul}(Z)$, and for $\mathfrak{e} \in \text{Eul}(Z)$ the maximal abelian torsion $\tau(Z, \mathfrak{e}, \omega)$ in a way completely analogous to the case of CW pairs. As in Definition 3.3, we also have a map $E_Z : \text{Lift}(Z) \rightarrow \text{Eul}(Z)$.

Remark. Notice that if Z is a nice handle decomposition of (M, γ) , then we can collapse each handle to its core, starting from e_n and proceeding to e_1 , and finally $S \times I$ to $S \times \{0\}$, to obtain a relative CW complex X built upon S . There are canonical bijections $\text{Lift}(Z) = \text{Lift}(X)$ and $\text{Eul}(Z) = \text{Eul}(X)$ such that given $\mathfrak{e} \in \text{Eul}(Z) = \text{Eul}(X)$ and a homology orientation ω , we have $\tau(Z, \mathfrak{e}, \omega) = \tau(X, \mathfrak{e}, \omega)$.

The sets $\text{Eul}(Z)$ and $\text{Spin}^c(M, \gamma)$ are both affine copies of $H_1(M)$. We will show that they are canonically isomorphic.

Claim. Let v_0 be a vector field as in subsection 2.2. Then there exists a vector field v_Z on M with the following properties: v_Z vanishes exactly at the centers of $d(e_1), \dots, d(e_n)$; the index of v_Z at the center of $d(e_i)$ is $(-1)^{I(i)}$; finally, $v_Z|_{\partial M} = v_0$.

Proof. We first construct such a vector field v_A on A , then push it forward using the diffeomorphism d . First, let $v_A|(S \times I) = \partial/\partial t$, where t is the coordinate on I . If $e_i = D^3$ is a 0-handle with coordinates (x, y, z) , then let

$$v_Z(x, y, z) = x \cdot \frac{\partial}{\partial x} + y \cdot \frac{\partial}{\partial y} + z \cdot \frac{\partial}{\partial z}.$$

On a 3-handle take v_A to be the negative of the previous vector field. If $e_j = D^1 \times D^2$ is a 1-handle with coordinates x on D^1 and (y, z) on D^2 , then we define $v_Z(x, y, z)$ to be

$$-x \cdot \frac{\partial}{\partial x} + y \cdot \frac{\partial}{\partial y} + z \cdot \frac{\partial}{\partial z}.$$

This vector field points out of e_j along $D^1 \times S^1$ and it lies on the same side of $S^0 \times D^2$ as $\{0\} \times D^2$. We can glue $v_A|(D^1 \times D^2)$ to $v_A|(S \times I) \cup (0\text{-handles})$ by a smooth handle attachment. Similarly, on a 2-handle $D^2 \times D^1$ with coordinates (x, y, z) we choose v_A to be $-x \cdot \partial/\partial x - y \cdot \partial/\partial y + z \cdot \partial/\partial z$. One can also think of v_Z as the gradient-like vector field for a Morse function compatible with the handle decomposition Z . This concludes the proof of the claim. $\square$

Definition 3.8. For $\mathfrak{e} \in \text{Eul}(Z)$ let $s_Z(\mathfrak{e}) \in \text{Spin}^c(M, \gamma)$ be the homology class of the vector field v that is obtained as follows. Pick an Euler chain θ representing $\mathfrak{e}$ that is a union of pairwise disjoint smoothly embedded arcs and circles inside $\text{Int}(M)$. Choose an open regular neighborhood $N(\theta)$ of θ . Then let $v = v_Z$ on $M \setminus N(\theta)$. If N_0 is a component of $N(\theta)$ diffeomorphic to B^3 , then extend v to N_0 as a nowhere zero vector field. This is possible since v_Z has an index 1 and an index -1 singularity inside N_0 . The homology class of v is independent of the choice of extension. If N_1 is a component of $N(\theta)$ diffeomorphic to $S^1 \times B^2$, then we get $v|_{N_1}$ from $v_Z|_{N_1}$ using Reeb tubularization, as described in [Tu90, p.639].

Note that we can avoid closed components of θ in the above definition except if (M, γ) is a product and $A = S \times I$.

Lemma 3.9. For $\mathfrak{e}_1, \mathfrak{e}_2 \in \text{Eul}(Z)$, we have $s_Z(\mathfrak{e}_1) - s_Z(\mathfrak{e}_2) = \mathfrak{e}_1 - \mathfrak{e}_2$.

Proof. An analogous obstruction theoretic argument as in the proof of [OS04a, Lemma 2.19] works here too. Also see [Tu90]. $\square$

Consequently, s_Z is an H -equivariant bijection between $\text{Eul}(Z)$ and $\text{Spin}^c(M, \gamma)$.

Remark. Note that if the handle decomposition Z arises from a balanced diagram (Σ, α, β) , then every $\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$ defines a unique Euler structure $\mathfrak{e}(\mathbf{x}) \in \text{Eul}(Z)$ as follows. Suppose that $x_i \in \alpha_j \cap \beta_k$, and let a_j be the 1-handle corresponding to α_j and b_k the 2-handle corresponding to β_k . Then let θ_i be a curve

that connects the center of a_j to x_i inside a_j and then goes from x_i to the center of b_k inside b_k . The Euler chain $\theta_1 + \dots + \theta_d$ defines $\mathfrak{e}(\mathbf{x})$. Then the Spin^c -structure $\mathfrak{s}(\mathbf{x})$ assigned to $\mathbf{x}$ is exactly $s_Z(\mathfrak{e}(\mathbf{x}))$.

Proposition 3.10. *Suppose that Z and Z' are nice handle decompositions of (M, γ) . Let $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ and choose a homology orientation ω of $(M, R_-(\gamma))$. Then*

$$\tau(Z, s_Z^{-1}(\mathfrak{s}), \omega) = \tau(Z', s_{Z'}^{-1}(\mathfrak{s}), \omega).$$

Proof. By [Ce70] one can get from Z to Z' through a finite sequence of nice handle decompositions, each one obtained from the previous by one of the following operations: a handle isotopy, a handle slide, adding/cancelling a pair of handles, or an isotopy of d . So it suffices to show the claim when $Z = (\mathcal{A}, d)$ and $Z' = (\mathcal{A}', d')$ are related by one of these operations.

Suppose that $\mathcal{A} = \mathcal{A}'$ and d is isotopic to d' . Then there is a diffeotopy $\{h_t: t \in I\}$ of M such that $d' = h_1 \circ d$. If the submanifold $\theta \subset M$ represents $s_Z^{-1}(\mathfrak{s})$, then $\theta' = d' \circ d^{-1}(\theta) = h_1(\theta)$ is isotopic to θ . Moreover, $v_{Z'} = d'_*(v_A) = (h_1)_*(v_Z)$ is isotopic to v_Z . Consequently, θ' represents $s_{Z'}^{-1}(\mathfrak{s})$ and both $\tau(Z, s_Z^{-1}(\mathfrak{s}), \omega)$ and $\tau(Z', s_{Z'}^{-1}(\mathfrak{s}), \omega)$ are equal to the torsion $\tau(A, S \times I, [d^{-1}(\theta)], \omega)$.

Now we consider the case when Z and Z' are related by isotoping a handle. This means the following. We choose a handle e_i of $\mathcal{A}$ and isotope its attaching map f_i to some other map f'_i inside S_{i-1} . Then we extend this isotopy to a diffeotopy $\{\varphi_t: t \in I\}$ of S_{i-1} such that $f'_i = \varphi_1 \circ f_i$. We define the A'_j recursively, together with diffeomorphisms $\nu_j: A_j \rightarrow A'_j$ for $j \geq i-1$. If $j \leq i-1$, then let $A'_j = A_j$. To define ν_{i-1} choose a collar $S_{i-1} \times I$ of S_{i-1} such that $S_{i-1} \times \{1\}$ is identified with S_{i-1} . Outside this collar let $\nu_{i-1} = \text{id}$, and $\nu_{i-1}(s, t) = (\varphi_t(s), t)$ for $(s, t) \in S_{i-1} \times I$. If A'_{j-1} and ν_{j-1} are already defined, then we obtain A'_j by gluing $e'_j = e_j$ to S'_{j-1} along $\nu_j \circ f_j$. Then ν_{j-1} naturally extends to A_j , call this extension ν_j . This defines the handle complex $\mathcal{A}'$, and we set $d' = d \circ (\nu_n)^{-1}$, where n is the number of handles.

Define $\nu = \nu_n$, this is a diffeomorphism from A to A' . For $1 \leq j \leq n$ let p_j be the center of e_j and p'_j the center of e'_j . Then $\nu(p_j) = p'_j$, so $d(p_j) = d'(p'_j)$. Hence there is a natural bijection $N: \text{Eul}(Z) \rightarrow \text{Eul}(Z')$ using the formula $N([\theta]) = [\theta]$. The vector field $\nu_*(v_A)$ agrees with $v_{A'}$ except on $\nu(S_{i-1} \times I)$. But there they both point up (with respect to $\partial/\partial t$), so $\nu_*(v_A)$ and $v_{A'}$ are isotopic on this collar through nowhere zero fields rel boundary. This proves that for every $\mathfrak{e} \in \text{Eul}(Z)$ we have $s_Z(\mathfrak{e}) = s_{Z'}(N(\mathfrak{e}))$.

Let $\hat{M}$ be the maximal abelian cover of M and let $\hat{Z}$ and $\hat{Z}'$ be the induced relative handle structures on $\hat{M}$. We show that $\tau(Z, \mathfrak{e}, \omega) = \tau(Z', N(\mathfrak{e}), \omega)$ for two special isotopies. Let θ be a submanifold simultaneously representing $\mathfrak{e}$ and $N(\mathfrak{e})$. We can assume that θ has no closed components. Let $\hat{\theta}$ be an arbitrary lift of θ to $\hat{M}$. Then $\partial\hat{\theta}$ defines a lift in both $\hat{Z}$ and $\hat{Z}'$ that represent $E_Z^{-1}(\mathfrak{e})$ and $E_{Z'}^{-1}(N(\mathfrak{e}))$, respectively. Let $\hat{e}_j$ be the lift of $d(e_j)$ and $\hat{e}'_j$ the lift of $d'(e'_j)$. Since the centers of the handles of $\hat{Z}$ and $\hat{Z}'$ coincide, we have a bijection $B: C_*(\hat{Z}) \rightarrow C_*(\hat{Z}')$ that takes the generator $\hat{e}_j$ to $\hat{e}'_j$.

First suppose that the isotopy connecting f_i and f'_i avoids every other handle of index $I(i)$, including the attaching map of e_j if $j > i$ and $I(j) = I(i)$. Then B is an isomorphism of based complexes. Indeed, isotoping the attaching map of $\hat{e}_i$ does not change the algebraic intersection number with belt circles of handles of index $I(i) - 1$.

The second case is when we handleslide e_i over a handle e_r such that $I(i) = I(r)$. Then we can obtain the based complex $C_*(\hat{Z}')$ from $C_*(\hat{Z})$ by replacing the basis element $\hat{e}_i$ with $\hat{e}_i + \hat{e}_r$. Thus the torsion is again unchanged.

Now suppose that Z' is obtained from Z by adding a canceling pair of handles e and f , and suppose that Z' is also nice (this is not necessarily the case if $I(i) = 1$, but that can be avoided by isotoping the 2-handles beforehand). Suppose that e and f are attached between e_{i-1} and e_i . Similarly to the case of an isotopy, we recursively define $\mathcal{A}'$ together with a sequence of diffeomorphisms $\nu_j: A_j \rightarrow A'_{j+2}$ for $j \geq i - 1$. Let $A'_j = A_j$ for $j \leq i - 1$. To define ν_{i-1} choose a collar $S_{i-1} \times I$ of S_{i-1} as before. Outside this collar let $\nu_{i-1} = \text{id}$, and $\nu_{i-1}(S_{i-1} \times I) = (S_{i-1} \times I) \cup e \cup f$. We set $e'_i = e$ and $e'_{i+1} = f$. If $j \geq i$ and the attaching map of e_j in $\mathcal{A}$ is f_j , then let $e'_{j+2} = e_j$ attached to A_{j-1} along $\nu_{j-1} \circ f_j$. Finally, set $\nu = \nu_n$ and define $d' = d \circ \nu^{-1}$.

The CW complexes corresponding to $\mathcal{A}$ and $\mathcal{A}'$ are related by an elementary expansion s , so by Lemma 3.4 there is a bijection $b_s: \text{Eul}(Z) \rightarrow \text{Eul}(Z')$ such that $\tau(Z, \mathbf{e}, \omega) = \tau(Z', b_s(\mathbf{e}), \omega)$ for every $\mathbf{e} \in \text{Eul}(Z)$. We claim that $s_Z(\mathbf{e}) = s_{Z'}(b_s(\mathbf{e}))$. Indeed, let p'_j denote the center of e'_j . Then if θ represents $\mathbf{e}$ and δ is an arc inside $e'_i \cup e'_{i+1}$ such that $\partial\delta = (-1)^i(p'_i - p'_{i+1})$, then $\theta' = \theta + d'(\delta)$ represents $b_s(\mathbf{e})$. Let $K = (S'_{i-1} \times I) \cup e'_i \cup e'_{i+1} \subset A'$, then $d'(K) = d(S_{i-1} \times I)$. If v' is a nowhere vanishing vector field on M extending $v_{Z'}|(M \setminus N(\theta'))$, then $v'|d'(K)$ is isotopic to $d_*(\partial/\partial t)$. On the other hand, if v is the nowhere zero vector field obtained from v_Z and θ , then $v|d'(K) = d_*(\partial/\partial t)$ and $v|(M \setminus d'(K)) = v'|(M \setminus d'(K))$. The claim follows. $\square$

Definition 3.11. Let Z be a nice handle decomposition of (M, γ) . Given an element $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ and a homology orientation ω for the pair $(M, R_-(\gamma))$, we define

$$\tau(M, \gamma, \mathfrak{s}, \omega) = \tau(Z, s_Z^{-1}(\mathfrak{s}), \omega) \in Q(H),$$

where s_Z^{-1} is the canonical identification between $\text{Spin}^c(M, \gamma)$ and $\text{Eul}(Z)$ defined in Definition 3.8. By Proposition 3.10 the torsion $\tau(M, \gamma, \mathfrak{s}, \omega)$ is independent of the choice of Z .

We now extend Definition 3.11 to disconnected weakly balanced sutured manifolds.

Definition 3.12. Suppose that (M, γ) is a weakly balanced sutured manifold whose components are $(M_1, \gamma_1), \dots, (M_n, \gamma_n)$. Fix a homology orientation ω for $(M, R_-(\gamma))$ and a Spin^c structure $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$. For $1 \leq i \leq n$ let $\mathfrak{s}_i = \mathfrak{s}|_{M_i}$ and $\omega_i = \omega|_{M_i}$. Then define

$$\tau(M, \gamma, \mathfrak{s}, \omega) = \bigotimes_{i=1}^n \tau(M_i, \gamma_i, \mathfrak{s}_i, \omega_i) \in \bigotimes_{i=1}^n Q(\mathbb{Z}[H_1(M_i)]) \subset Q(H),$$

where we take the tensor product over $\mathbb{Z}$.

Now suppose that $R_+(\gamma) \cap M_i \neq \emptyset$ and $R_-(\gamma) \cap M_i \neq \emptyset$ for every $1 \leq i \leq n$. This is true for example if (M, γ) is balanced. Then (M, γ) has a handle decomposition with no 0 and 3–handles. From Lemma 3.5 we get that $\tau(M, \gamma, \mathfrak{s}, \omega) \in \mathbb{Z}[H]$. Following Turaev, we define the torsion function as

$$\begin{aligned} T_{(M, \gamma, \omega)} : \text{Spin}^c(M, \gamma) &\rightarrow \mathbb{Z} \\ \mathfrak{s} &\mapsto \tau(M, \gamma, \mathfrak{s}, \omega)_1 \end{aligned}$$

where $\tau(M, \gamma, \mathfrak{s}, \omega)_1$ denotes the constant term of $\tau(M, \gamma, \mathfrak{s}, \omega) \in \mathbb{Z}[H]$. Note that in light of (2) we can recover $\tau(M, \gamma, \mathfrak{s}, \omega) \in \mathbb{Z}[H]$ from the function $T_{(M, \gamma, \omega)}$.

3.5. Making γ connected. Let (M, γ) be a connected balanced sutured manifold and fix a homology orientation ω . In the future, it will often be convenient to assume that γ , and hence $R_\pm(\gamma)$ are connected. This can be arranged by adding product 1–handles to (M, γ) to produce a new sutured manifold (M', γ') . In this section, we describe the effect of this operation on the sutured Floer homology and the torsion. In particular, we show that $SFH(M, \gamma)$ can be recovered from $SFH(M', \gamma')$, and likewise for the torsion.

In terms of Heegaard diagrams, this operation of adding a product 1–handle can be described as follows. Suppose $(\Sigma, \boldsymbol{\alpha}, \boldsymbol{\beta})$ is a balanced sutured Heegaard diagram for (M, γ) , and let Σ' be the result of attaching a 2–dimensional 1–handle h to Σ . Then $(\Sigma', \boldsymbol{\alpha}, \boldsymbol{\beta})$ is a Heegaard diagram representing a sutured manifold (M', γ') , which is obtained from M by attaching the 3–dimensional 1–handle $h \times [-1, 1]$. We can recover (M, γ) from (M', γ') by decomposing along the product disk $c \times [-1, 1]$, where c is the cocore of h . If h joins two different components of γ , then γ' will have one less component than γ .

The embedding $(M, R_-(\gamma)) \hookrightarrow (M', R_-(\gamma'))$ induces an isomorphism

$$H_*(M, R_-(\gamma); \mathbb{R}) \rightarrow H_*(M', R_-(\gamma'); \mathbb{R}).$$

Hence ω canonically gives rise to an orientation ω' of $H_*(M', R_-(\gamma'); \mathbb{R})$.

From [Ju06, Lemma 9.13] we know that $SFH(M', \gamma') \cong SFH(M, \gamma)$. To be more precise, by [Ju08b, Prop. 5.4] there is an injection $i : \text{Spin}^c(M, \gamma) \rightarrow \text{Spin}^c(M', \gamma')$ for which

$$SFH(M, \gamma, \mathfrak{s}) \cong SFH(M', \gamma', i(\mathfrak{s})).$$

Moreover, $SFH(M', \gamma', \mathfrak{s}') = 0$ if $\mathfrak{s}'$ is not in the image of i . The same proof shows that using the homology orientations ω and ω' we get an isomorphism of $\mathbb{Z}/2$ graded groups.

The injection i is most easily described by observing that the generating sets $\mathbb{T}_\alpha \cap \mathbb{T}_\beta$ and $\mathbb{T}_{\alpha'} \cap \mathbb{T}_{\beta'}$ are naturally identified, and setting $i(\mathfrak{s}(\mathbf{x})) = \mathfrak{s}(\mathbf{x}')$, where $\mathbf{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$ and $\mathbf{x}' \in \mathbb{T}_{\alpha'} \cap \mathbb{T}_{\beta'}$ are corresponding intersection points. This map extends to a map on $\text{Spin}^c(M, \gamma)$ using the free and transitive $H_1(M)$ –action, which extends to a free action on $\text{Spin}^c(M', \gamma')$ using the injection $H_1(M) \rightarrow H_1(M')$.

More intrinsically, i can be defined as follows. Suppose that $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ is represented by some vector field v . By definition, $v|_\gamma$ is the gradient of a height function $\gamma \rightarrow [-1, 1]$. So we get a smooth vector field v' on M' if we let $v'|_M = v$ and $v'|_h = p^*(\partial/\partial t)$, where $p: h \times [-1, 1] \rightarrow [-1, 1]$ is the projection. We define $i(\mathfrak{s})$ as the homology class of v' .

To see that the latter description coincides with the former, let Z and Z' be the handle decompositions arising from (Σ, α, β) and (Σ', α, β) , respectively. The inclusion $M \hookrightarrow M'$ naturally induces an affine injection $i_0: \text{Eul}(Z) \rightarrow \text{Eul}(Z')$, and i is the composition $s_{Z'} \circ i_0 \circ s_Z^{-1}$. It is easy to see that $i_0(\mathfrak{e}(\mathbf{x})) = \mathfrak{e}(\mathbf{x}')$, and this implies that $i(\mathfrak{s}(\mathbf{x})) = \mathfrak{s}(\mathbf{x}')$.

An analogous statement is satisfied by the torsion:

Lemma 3.13. *Suppose that (M', γ') is obtained from the connected balanced sutured manifold (M, γ) by adding a product 1-handle. Then $T_{(M, \gamma, \omega)}(\mathfrak{s}) = T_{(M', \gamma', \omega')}(i(\mathfrak{s}))$ for every $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$. Moreover, $T_{(M', \gamma', \omega')}(\mathfrak{s}') = 0$ if $\mathfrak{s}'$ is not in the image of i .*

Proof. Let Z be the handle decomposition given by (Σ, α, β) and Z' the decomposition given by (Σ', α, β) . The universal abelian cover $p': \hat{M}' \rightarrow M$ can be constructed as follows. Start with a disjoint union $\sqcup_{i \in \mathbb{Z}} \hat{M}_i$, where each $\hat{M}_i$ is homeomorphic to the universal abelian cover $\hat{M}$ of M . Now join $\hat{M}_i$ to $\hat{M}_{i+1}$ by 1-handles, one for each element of $H_1(M)$. These 1-handles are all thickenings of 1-handles in $\hat{R}'_- = (p')^{-1}(R_-(\gamma'))$, so they do not contribute to $C_*(\hat{M}', \hat{R}'_-)$. Choose a basis for $C_*(\hat{M}', \hat{R}'_-)$ all of whose handles are contained in $\hat{M}_0$; let $\mathfrak{l}$ and $\mathfrak{l}'$ be the associated lifts for M and M' . With respect to such a basis,

$$C_*(\hat{M}', \hat{R}'_-) \cong C_*(\hat{M}, \hat{R}_-) \otimes_{\mathbb{Z}} \mathbb{Z}[t, t^{-1}].$$

It follows that $\tau(Z', \mathfrak{l}', \omega') = i_*(\tau(Z, \mathfrak{l}, \omega))$, where $i_*: H_1(M) \rightarrow H_1(M')$ is the inclusion. It is easy to see that $i_0(E_Z(\mathfrak{l})) = E_{Z'}(\mathfrak{l}')$. Since $i = s_{Z'} \circ i_0 \circ s_Z^{-1}$, it follows that $\tau(M', \gamma', i(\mathfrak{s}), \omega') = i_*(\tau(M, \gamma, \mathfrak{s}, \omega))$. It is now straightforward to see that this implies the statement of the lemma. $\square$

4. THE PROOF OF THEOREM 1

We are now in a position to prove that the Euler characteristic of $SFH(M, \gamma, \mathfrak{s}, \omega)$ coincides with the torsion $T_{(M, \gamma, \omega)}(\mathfrak{s})$. Before giving the proof, we recall some basic facts about handle decompositions, presentations of π_1 , and Fox calculus.

4.1. Balanced diagrams and presentations. We begin by explaining how to find a presentation of $\pi_1(M)$ compatible with a nice handle decomposition Z having no 0 and 3-handles, or equivalently, with a balanced diagram (Σ, α, β) representing (M, γ) . Choose a 2-dimensional handle decomposition of R_- consisting of one 0-handle and l 1-handles; this naturally gives a 3-dimensional handle decomposition of $R_- \times I$, again with one 0-handle and l 1-handles. Without loss of generality, we may assume that the attaching disks of the 1-handles of Z are disjoint from the belt circles of the

1-handles of $R_- \times I$, and thus (after an isotopy) that the 1-handles of Z are attached to the 0-handle of $R_- \times I$.

Fix a basepoint $p \in R_- \times \{0\}$ which is contained in the 0-handle. Then $\pi_1(M, p)$ is generated by loops $\alpha_1^*, \dots, \alpha_d^*, c_1^*, \dots, c_l^*$, where α_i^* runs through the i -th 1-handle of Z , and c_k^* runs through the k -th 1-handle of $R_- \times I$. (Note that α_i^* depends on the choice of handle decomposition for R_- as well as on Z .) Each 2-handle in Z gives rise to a relation as follows. Choose a path q_j from p to the attaching circle β_j of the j -th 2-handle; then $\bar{\beta}_j = q_j \beta_j q_j^{-1}$ is a loop which represents a trivial element of $\pi_1(M, p)$. We have a presentation

$$\pi_1(M, p) = \langle \alpha_1^*, \dots, \alpha_d^*, c_1^*, \dots, c_l^* \mid \bar{\beta}_1, \dots, \bar{\beta}_d \rangle.$$

To read off this presentation from a sutured Heegaard diagram (Σ, α, β) compatible with Z , we proceed as follows. First, surger S along the α -curves to produce a surface homeomorphic to R_- and containing $2d$ marked disks (the traces of the surgery). Next, choose a system of disjoint properly embedded arcs $c_1, \dots, c_l$ in R_- whose complement is homeomorphic to a disk. (This amounts to choosing a handlebody decomposition of R_- .) Without loss of generality, we may assume that the c_k 's are disjoint from the marked disks, so they lift to arcs $c_1, \dots, c_l$ in Σ . To write down the word $\bar{\beta}_j$, we simply traverse $\bar{\beta}_j$ and record its intersections with the α_i 's and the c_k 's as we go.

More precisely, an intersection point x between α_i and $\bar{\beta}_j$ is recorded by α_i^* if the sign of intersection $\alpha_i \cdot \bar{\beta}_j$ at x is positive, and by $(\alpha_i^*)^{-1}$ if the sign of the intersection is negative. Note that in doing this, we have implicitly chosen orientations on the α_i 's, the β_j 's, and the c_k 's.

In what follows, it will be convenient to choose the path q_j such that its image in Σ does not intersect any of the α_i or c_k . (This is always possible, since the complement of the α 's and c 's is connected.) In this case, we write the resulting relation as β_j ; it is obtained by traversing the curve β_j and recording the intersections with the α 's and the c 's.

4.2. Torsion from a balanced diagram. Given a nice handle decomposition of (M, γ) as in the previous section, the torsion $\tau(M, \gamma)$ can be computed as follows. The based curves $c_1^*, \dots, c_l^*$ give rise to a basis $\hat{C}_1^*, \dots, \hat{C}_l^*$ of $C_1(R_-; \mathbb{Z}[H])$. Similarly, the based curves $\alpha_1^*, \dots, \alpha_d^*$ give a basis $\hat{A}_1^*, \dots, \hat{A}_d^*$ of $C_1(M, R_-; \mathbb{Z}[H])$. Let B_j denote the 2-handle attached along β_j . Then the basings $q_1, \dots, q_n$ give rise to a choice of lifts $\hat{B}_1, \dots, \hat{B}_d$ of $B_1, \dots, B_d$. These give a basis for $C_2(M; \mathbb{Z}[H])$.

Proposition 4.1. *Let $\mathfrak{l} \in \text{Lift}(Z)$ be the element corresponding to the basings. Then up to sign we have*

$$\tau(M, \gamma, \mathfrak{l}) = \det \left(\varphi \left(\frac{\partial \bar{\beta}_j}{\partial \alpha_i^*} \right) \right) \in \mathbb{Z}[H].$$

Proof. Consider the following diagram of free $\mathbb{Z}[H]$ -modules. We write our choice of basis under the free modules.

$$\begin{array}{ccccccc}
& 0 & \rightarrow & C_2(M; \mathbb{Z}[H]) & \rightarrow & C_2(M, R_-; \mathbb{Z}[H]) & \rightarrow 0 \\
& & & \hat{B}_1, \dots, \hat{B}_d & & \hat{B}_1, \dots, \hat{B}_d & \\
& & \downarrow & \downarrow & & \downarrow & \\
0 & \rightarrow & C_1(R_-; \mathbb{Z}[H]) & \rightarrow & C_1(M; \mathbb{Z}[H]) & \rightarrow & C_1(M, R_-; \mathbb{Z}[H]) \rightarrow 0 \\
& & \hat{C}_1^*, \dots, \hat{C}_l^* & \rightarrow & \hat{A}_1^*, \dots, \hat{A}_d^* & & \hat{A}_1^*, \dots, \hat{A}_d^* \\
& & \downarrow & \downarrow & & \downarrow & \\
0 & \rightarrow & C_0(R_-; \mathbb{Z}[H]) & \rightarrow & C_0(M; \mathbb{Z}[H]) & \rightarrow & 0. \\
& & \hat{p} & & \hat{p} & &
\end{array}$$

Fox calculus tells us that the boundary map $C_2(M; \mathbb{Z}[H]) \rightarrow C_1(M; \mathbb{Z}[H])$ is given by

$$\begin{pmatrix} \varphi \left(\frac{\partial \bar{\beta}_j}{\partial \alpha_i^*} \right) \\ \varphi \left(\frac{\partial \bar{\beta}_j}{\partial c_k^*} \right) \end{pmatrix}.$$

The proposition now follows immediately. $\square$

4.3. Equality in $\mathbb{Z}[H]/\pm H$. We now turn to the proof of Theorem 1. First, observe that it is enough to prove the equality in the case where $R_-(\gamma)$ is connected. Indeed, if $R_-(\gamma)$ is not connected, then we can add product one-handles to obtain a new sutured manifold (M', γ') with $R_-(\gamma')$ connected. Lemma 3.13 and the discussion preceding it show that if $\chi(SFH(M', \gamma', i(\mathfrak{s}), \omega')) = T_{(M', \gamma', \omega')}(i(\mathfrak{s}))$, then $\chi(SFH(M, \gamma, \mathfrak{s}, \omega)) = T_{(M, \gamma, \omega)}(\mathfrak{s})$.

For the rest of this section, we assume $R_-(\gamma)$ is connected. Our next step is to show that the torsion and the Euler characteristic agree up to multiplication by $\pm[h]$ for some $h \in H$. In light of Proposition 4.1, it suffices to prove the following.

Proposition 4.2.

$$\chi(SFH(M, \gamma)) \sim \det \varphi \left(\frac{\partial \beta_j}{\partial \alpha_i^*} \right),$$

where $\varphi : \mathbb{Z}[\pi_1(M)] \rightarrow \mathbb{Z}[H_1(M)]$ is the homomorphism induced by abelianization and where $\sim$ indicates equality up to multiplication by $\pm[h]$.

Proof. We argue along the lines of chapter 3 in [Ra03]. First, observe that there are natural bijections

$$\{\text{elements of } \alpha_i \cap \beta_j\} \leftrightarrow \{\text{appearances of } \alpha_i^* \text{ in } \beta_j\} \leftrightarrow \left\{ \text{monomials in } \varphi \left(\frac{\partial \beta_j}{\partial \alpha_i^*} \right) \right\},$$

where the free derivative $\partial\beta_j/\partial\alpha_i^*$ has been expanded *without canceling any terms*. Equivalently,

$$\varphi\left(\frac{\partial\beta_j}{\partial\alpha_i^*}\right) = \sum_{x \in \alpha_i \cap \beta_j} m(x)[A(x)],$$

where each $m(x) = \pm 1$ and each $A(x)$ is an element of $H_1(M)$. The sign $m(x)$ is given by the exponent of the corresponding appearance of α_i^* in β_j , or equivalently, by the sign of intersection $\alpha_i \cdot \beta_j$ at x .

Recall that the chain complex computing SFH is generated by d -tuples of intersection points $\mathbf{x} = \{x_1, \dots, x_d\}$, where $x_i \in \alpha_i \cap \beta_{\sigma(i)}$ for some permutation $\sigma \in S_d$. On the other hand, the determinant of a $d \times d$ matrix B_{ij} can be expanded as

$$\det B_{ij} = \sum_{\sigma \in S_d} \text{sign}(\sigma) B_{1\sigma(1)} \cdots B_{d\sigma(d)}.$$

Thus we get a bijection

$$\mathbb{T}_\alpha \cap \mathbb{T}_\beta \longleftrightarrow \left\{ \text{monomials in } \det \varphi\left(\frac{\partial\beta_j}{\partial\alpha_i^*}\right) \right\}.$$

Again, all terms in the determinant are to be expanded *without cancellation*.

Together with Lemma 2.8, these imply that

$$\tau(M, \gamma) \sim \det \varphi\left(\frac{\partial\beta_j}{\partial\alpha_i^*}\right) = \sum_{\mathbf{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta} m(\mathbf{x})[A(\mathbf{x})].$$

Here $A(\mathbf{x}) = \sum_{i=1}^d A(x_i)$ and $m(\mathbf{x})$ is the sign of the intersection $\mathbb{T}_\alpha \cdot \mathbb{T}_\beta$ at $\mathbf{x}$. The orientations on $\mathbb{T}_\alpha$ and $\mathbb{T}_\beta$ are induced by the orderings $\langle \alpha_1, \dots, \alpha_d \rangle$ and $\langle \beta_1, \dots, \beta_d \rangle$. On the other hand, we know from equation 1 at the end of section 2.5 that

$$\chi(SFH(M, \gamma)) \sim \sum_{\mathbf{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta} m(\mathbf{x})[\iota(\mathfrak{s}(\mathbf{x}))],$$

where $\iota : \text{Spin}^c(M, \gamma) \rightarrow H_1(M)$ is an affine isomorphism. Comparing, we see that the two expressions will agree up to multiplication by $\pm[h]$ if we can show the following.

Lemma 4.3. $A(\mathbf{x}) - A(\mathbf{y}) = \mathbf{x} - \mathbf{y}$.

Proof. The coefficients $A(x)$ appearing in the Fox derivative $\partial\beta_j/\partial\alpha_i^*$ can be interpreted geometrically as follows. In the universal abelian cover $\hat{M}$, we have fixed lifts $\hat{\alpha}_i$ of α_i and $\hat{\beta}_j$ of β_j coming from the basings. An intersection point $x \in \alpha_i \cap \beta_j$ lifts to a unique $\hat{x} \in \hat{\beta}_j$, and this point $\hat{x}$ is contained in $(A(x) \cdot \hat{\alpha}_i) \cap \hat{\beta}_j$.

Suppose $\mathbf{x} = \{x_1, \dots, x_d\}$ and $\mathbf{y} = \{y_1, \dots, y_d\}$, and that the ordering of the x 's and y 's is arranged such that $x_j, y_j \in \beta_j$. Then there are i and i' such that $\hat{x}_j \in (A(x_j) \cdot \hat{\alpha}_i) \cap \hat{\beta}_j$ and $\hat{y}_j \in (A(y_j) \cdot \hat{\alpha}_{i'}) \cap \hat{\beta}_j$. If $\tilde{\beta}_j$ is an arbitrary lift of β_j , and

$\tilde{x}_j, \tilde{y}_j$ are the lifts of x_j, y_j which lie on it, then we can write $\tilde{x}_j \in (\tilde{A}(x_j) \cdot \hat{\alpha}_i) \cap \tilde{\beta}_j$ and $\tilde{y}_j \in (\tilde{A}(y_j) \cdot \hat{\alpha}_{i'}) \cap \tilde{\beta}_j$. Then we have

$$\tilde{A}(x_j) - \tilde{A}(y_j) = A(x_j) - A(y_j).$$

Now suppose that we are given a 1-cycle θ which runs from $\mathbf{x}$ to $\mathbf{y}$ along the α curves and a 1-cycle η which runs from $\mathbf{x}$ to $\mathbf{y}$ along the β curves. The closed 1-cycle $\theta - \eta$ may be divided into components. Let δ be one such component. If we start at some y_{j_1} on δ , and traverse δ , we will successively encounter points labeled $x_{j_1}, y_{j_2}, x_{j_2}, \dots, y_{j_r}, x_{j_r}$ before returning to y_{j_1} .

Let $\hat{\delta}$ be the lift of δ to $\hat{M}$ which starts at $\hat{y}_{j_1}$ and ends at $[\delta] \cdot \hat{y}_{j_1}$. As we traverse $\hat{\delta}$, we denote the α curve containing the lift of y_{j_k} by $\tilde{A}(y_{j_k}) \cdot \hat{\alpha}_{i'_k}$. Similarly, we denote the α curve containing the lift of x_{j_k} by $\tilde{A}(x_{j_k}) \cdot \hat{\alpha}_{i_k}$. Since x_{j_k} is joined to $y_{j_{k+1}}$ along an α curve, $\tilde{A}(x_{j_k}) = \tilde{A}(y_{j_{k+1}})$. On the other hand, since y_{j_k} is joined to x_{j_k} by a β curve, we have

$$\tilde{A}(x_{j_k}) - \tilde{A}(y_{j_k}) = A(x_{j_k}) - A(y_{j_k}).$$

We now compute

$$\begin{aligned} [\delta] &= \tilde{A}(x_{j_r}) - \tilde{A}(y_{j_1}) \\ &= \sum_{k=1}^r \left(\tilde{A}(x_{j_k}) - \tilde{A}(y_{j_k}) \right) \\ &= \sum_{k=1}^r (A(x_{j_k}) - A(y_{j_k})). \end{aligned}$$

A similar relation holds for each component of $\theta - \eta$. Adding them all together, we find that

$$\mathbf{x} - \mathbf{y} = [\theta - \eta] = \sum_{i=1}^n (A(x_i) - A(y_i)) = A(\mathbf{x}) - A(\mathbf{y}).$$

□

This completes the proof of Proposition 4.2

□

4.4. Spin^c structures. Our next step is to show that $\chi(SFH(M, \gamma, \mathfrak{s})) = \pm T_{(M, \gamma)}(\mathfrak{s})$ for every $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$. To see this, fix a single generator $\mathbf{x} = \{x_1, \dots, x_d\} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$ and consider the associated lift $\mathfrak{l}(\mathbf{x}) = E_Z^{-1}(\mathfrak{e}(\mathbf{x})) \in \text{Lift}(M, R_-(\gamma))$. We will show that the term in

$$\det \varphi \left(\frac{\partial \beta_j}{\partial \alpha_i^*} \right)$$

corresponding to $\mathbf{x}$ contributes to $T_{(M, \gamma)}(\mathfrak{s}(\mathbf{x}))$, or equivalently, that it contributes ± 1 to the torsion $\tau(M, \gamma, \mathfrak{l}(\mathbf{x}))$. It will then follow from Proposition 4.2 that the same relation holds for every generator $\mathbf{y} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$.

Without loss of generality, we may assume $x_i \in \alpha_i \cap \beta_i$. Consider the matrix

$$F_{ij} = \varphi \left(\frac{\partial \beta_j}{\partial \alpha_i^*} \right)$$

of Fox derivatives. Fix lifts $\tilde{A}_i^*$ of the 1-handles to the universal abelian cover. Then after an appropriate normalization (multiplying each column in the matrix by a unit in $\mathbb{Z}[H_1(M)]$), the column vector

$$\left(\varphi \left(\frac{\partial \beta_j}{\partial \alpha_1^*} \right), \dots, \varphi \left(\frac{\partial \beta_j}{\partial \alpha_d^*} \right) \right)^T$$

expresses the boundary of the lift $\tilde{B}_j$ of the two-handle B_j in terms of the lifts $\tilde{A}_i^*$. It is well-defined up to multiplication by a unit in $\mathbb{Z}[H_1(M)]$, corresponding to changing the lift $\tilde{B}_j$. Let us choose the $\tilde{B}_j$ such that the monomial $A(x_i)$ in F_{ii} is ± 1 . This choice of basis is clearly compatible with the Euler structure $\mathfrak{e}(\mathbf{x})$, so the corresponding element of $\text{Lift}(Z)$ is $\mathfrak{l}(\mathbf{x})$. Then

$$\tau(M, \gamma, \mathfrak{l}(\mathbf{x})) = \det \varphi \left(\frac{\partial \beta_j}{\partial \alpha_i^*} \right),$$

and the monomial $A(\mathbf{x})$ contributes to $\tau(M, \gamma, \mathfrak{l}(\mathbf{x}))$ with coefficient ± 1 . In other words, it is assigned to the Spin^c structure $\mathfrak{s}(\mathbf{x})$.

4.5. Homology orientations. To complete the proof of Theorem 1, it remains to check that if we fix a homology orientation ω for $H_*(M, R_-(\gamma); \mathbb{R})$, then we have $\chi(SFH(M, \gamma, \mathfrak{s}, \omega)) = T_{(M, \gamma, \omega)}(\mathfrak{s})$. This follows easily from the definitions. To compute the torsion, we choose bases $A_1^*, \dots, A_d^*$ for $C_1(Z)$ and $B_1, \dots, B_d$ for $C_2(Z)$ which induce ω ; we lift these bases to bases of $C_*(\tilde{Z})$, and the torsion is given by the determinant of the boundary map $\partial : C_2(\tilde{Z}) \rightarrow C_1(\tilde{Z})$, or equivalently, of $F = \varphi \left(\frac{\partial \beta_j}{\partial \alpha_i^*} \right)$.

To fix an absolute $\mathbb{Z}/2$ grading on $SFH(M, \gamma)$, we must orient the tori $\mathbb{T}_\alpha$ and $\mathbb{T}_\beta$. By Lemma 2.6, this choice determines a homology orientation ω for $H_*(M, R_-(\gamma); \mathbb{R})$. With respect to ω , the sign with which a generator $\mathbf{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$ contributes to the Euler characteristic is the intersection sign $\mathbb{T}_\alpha \cdot \mathbb{T}_\beta$ at $\mathbf{x}$. But this sign is precisely the sign of the corresponding term in $\det F$. $\square$

5. THE TORSION FUNCTION VIA FOX CALCULUS

In this section, we explain how to compute $T_{(M, \gamma)}$ using Fox calculus. We assume throughout that (M, γ) is balanced and that the subsurfaces $R_\pm(\gamma)$ are connected. In light of Lemma 3.13, this restriction is a very mild one. For brevity, we write $R_\pm$ for $R_\pm(\gamma)$ and H for $H_1(M)$.

Suppose we are given a basepoint $p \in R_-$ and a presentation

$$\pi_1(M, p) \cong \langle a_1, \dots, a_m \mid b_1, \dots, b_n \rangle.$$

We say that the presentation is *geometrically balanced* if $m - n = g(\partial M)$. For example, any presentation coming from a handle decomposition of M is geometrically balanced.

We now consider the map $e_* : \pi_1(R_-, p) \rightarrow \pi_1(M, p)$ induced by the embedding $e : R_- \rightarrow M$. Since R_- is connected and has nonempty boundary, $\pi_1(R_-, p)$ is free. Choose elements $d_1, \dots, d_l$ which freely generate $\pi_1(R_-, p)$, and let $e_k = e_*(d_k)$ be their images in $\pi_1(M, p)$ for $1 \leq k \leq l$. Expressing each e_k as a word in the a_i , we have the Fox derivatives $\frac{\partial e_k}{\partial a_i} \in \mathbb{Z}[\pi_1(M)]$. Denote by $\varphi : \mathbb{Z}[\pi_1(M)] \rightarrow \mathbb{Z}[H_1(M)]$ the homomorphism induced by abelianization. Then we can form the matrix

$$A = \left(\varphi \left(\frac{\partial b_j}{\partial a_i} \right) \quad \varphi \left(\frac{\partial e_k}{\partial a_i} \right) \right).$$

If our presentation of $\pi_1(M, p)$ and the sutured manifold (M, γ) are both balanced, then $1 - l = \chi(R_-) = 1 - g(\partial M) = 1 - m + n$, so $m = n + l$, and A is a square matrix. The main result of this section is the following proposition.

Proposition 5.1. *Let (M, γ) be a balanced sutured manifold such that M is irreducible. If A is a square matrix, then*

$$\tau(M, \gamma) = \det(A)$$

up to multiplication by an element in $\pm H$.

Remark. The above proposition was already proved by Goda and Sakasai in [GS08, Proposition 4.6] when (M, γ) is a homology cylinder.

Proof. If $M = D^3$, then the proposition is obviously true. Otherwise, since M is irreducible, no component of ∂M can be homeomorphic to S^2 , thus $b_1(M) > 0$.

As in Section 4.1, pick a nice handle decomposition Z for (M, γ) , and another handle decomposition for R_- . Again, denote by $c_1^*, \dots, c_l^*$ the corresponding curves in R_- based at a point p . Let Y be the 1-complex given by the 0-cell p and the 1-cells $c_1^*, \dots, c_l^*$. Let X be the 2-complex given by collapsing the handles of Z to cells. Note that X is homotopy equivalent to M and that Y is a subcomplex of X . Furthermore, $\tau(X, Y) = \tau(M, \gamma)$, cf. the remark after Definition 3.7.

Now let $\Pi = \langle a_1, \dots, a_m \mid b_1, \dots, b_n \rangle$ be a geometrically balanced presentation of $\pi_1(M, p)$, and let $d_1, \dots, d_l$ be any basis for the free group $\pi_1(R_-)$. A classical theorem of Nielsen in [Ni24] on automorphisms of free groups implies that any two bases $\langle e_1, \dots, e_l \rangle$ and $\langle f_1, \dots, f_l \rangle$ of a free group $\pi_1(R_-)$ are related by a sequence of the following moves. We either 1) replace e_k by $(e_k)^{-1}$ or 2) replace e_k by $e_k e_{k'}$ for some $k \neq k'$. Move 1) simply multiplies the k -th column of A by $-\varphi(e_k^{-1})$, while move 2) adds $\varphi(e_{k'})$ times the k' -th column to the k -th column. In both cases, the determinant of A changes only by multiplication by an element of $\pm[H]$. Thus we may assume that $d_k = [c_k^*]$ for $k = 1, \dots, l$.

Let D be the 2-complex with one 0-cell p and 1-cells $A_1, \dots, A_m$ corresponding to $a_1, \dots, a_m$, finally 2-cells $B_1, \dots, B_n$ glued along $b_1, \dots, b_n$, respectively. We denote by X' the 2-complex obtained from D by adding extra 1-cells $C_1^*, \dots, C_l^*$ corresponding

to $c_1^*, \dots, c_l^*$, and by adding 2-cells $W_1, \dots, W_l$ such that ∂W_i is glued along $e_*([c_i^*]) \cdot (C_i^*)^{-1}$. Note that X' is a 2-complex that is simple homotopy equivalent to D . We can and will view Y as a subcomplex of X' .

We have an obvious isomorphism $\psi : \pi_1(X') \rightarrow \pi_1(X)$ which induces the identity on $\pi_1(Y)$, viewed as a subgroup of both $\pi_1(X')$ and $\pi_1(X)$. Since M is irreducible and $\pi_1(M)$ is infinite, M , and hence X , is aspherical. It follows that there exists a map $f : X' \rightarrow X$ with $f_* = \psi$ and such that $f|_Y = \text{id}_Y$.

Claim. The map $f : X' \rightarrow X$ induces an isomorphism $H_2(X') \rightarrow H_2(X)$.

Let F be the homotopy fiber corresponding to the map $f : X' \rightarrow X$. Note that $\pi_1(F) = 0$, since $f_* : \pi_1(X') \rightarrow \pi_1(X)$ is an isomorphism and $\pi_2(X) = 0$. It follows that $H_1(F) = 0$. Now consider the Serre spectral sequence corresponding to the fibration $F \rightarrow X' \rightarrow X$: We have a spectral sequence with $E_{p,q}^2 = H_p(X; H_q(F))$ which converges to $H_*(X')$. Note that $E_{p,q}^2 = 0$ for $p > 2$ since X is a 2-complex, and we also have $E_{p,1}^2 = 0$ since $H_1(F) = 0$. It follows that there is a short exact sequence

$$0 \rightarrow H_0(X; H_2(F)) \rightarrow H_2(X') \rightarrow H_2(X) \rightarrow 0.$$

In particular, the map $H_2(X') \rightarrow H_2(X)$ is surjective. Since X and X' are 2-complexes, both $H_2(X)$ and $H_2(X')$ are free abelian groups. We also have $H_i(X') \cong H_i(X)$ for $i = 0, 1$, and

$$\chi(X') = 1 - (m + l) + n + l = 1 - m + n = \chi(R_-) = \chi(M) = \chi(X).$$

Hence the map $H_2(X') \rightarrow H_2(X)$ is an epimorphism between torsion free abelian groups of the same rank, so it is an isomorphism.

Claim. The map $f : X' \rightarrow X$ induces a homotopy equivalence of pairs $(X', Y) \rightarrow (X, Y)$.

First note that $f_* : \pi_1(X') \rightarrow \pi_1(X)$ is an isomorphism. We write $\pi = \pi_1(X)$ and identify $\pi_1(X')$ with π via f_* . Recall that $\pi_i(X; \mathbb{Z}[\pi]) = 0$ for $i > 1$ since X is aspherical. This implies that $H_i(X; \mathbb{Z}[\pi]) = 0$ for $i > 1$, so it suffices to show that $H_i(X'; \mathbb{Z}[\pi]) = 0$ for all $i > 1$.

Let U be the mapping cylinder of $f : X' \rightarrow X$. Then U is a 3-complex which is homotopy equivalent to X and contains X' . We can identify π with $\pi_1(U)$. From the long exact sequence of the pair (U, X') we now obtain that $H_2(X'; \mathbb{Z}[\pi]) \cong H_3(U, X'; \mathbb{Z}[\pi])$. So it remains to show that $H_3(U, X'; \mathbb{Z}[\pi]) = 0$.

By the previous claim, the map $f_* : H_2(X') \rightarrow H_2(U) = H_2(X)$ is an isomorphism. It follows that $H_3(U, X') = 0$. In particular, the map of free $\mathbb{Z}[\pi]$ -modules $C_3(U, X'; \mathbb{Z}[\pi]) \rightarrow C_2(U, X'; \mathbb{Z}[\pi])$ becomes injective after tensoring over $\mathbb{Z}[\pi]$ with $\mathbb{Z}$. On the other hand, by [Ho82, Corollary 6.2] the group π is locally indicable. According to [HS83, Theorem 1], this implies that $C_3(U, X'; \mathbb{Z}[\pi]) \rightarrow C_2(U, X'; \mathbb{Z}[\pi])$ is injective, and in particular $H_3(U, X'; \mathbb{Z}[\pi])$ is trivial. This concludes the proof of the claim.

By hypothesis, M is irreducible and $b_1(M) > 0$, so M is Haken. By a theorem of Waldhausen [Wa78], the Whitehead group of $\pi = \pi_1(M)$ vanishes. Thus the map $f : (X', Y) \rightarrow (X, Y)$ is a simple homotopy equivalence, and the maximal abelian torsions of (X', Y) and (X, Y) agree. We have now shown that

$$\tau(M, \gamma) = \tau(X, Y) = \tau(X', Y).$$

The following claim therefore concludes the proof of the proposition.

Claim. We have $\tau(X', Y) = \det(A)$.

As before, we pick lifts of the cells in (X', Y) to the universal abelian cover. These lifts are, as usual, decorated by a ‘hat’. Consider the following diagram of free $\mathbb{Z}[H]$ -modules.

$$\begin{array}{ccccccc}
0 & & \rightarrow & C_2(X'; \mathbb{Z}[H]) & \rightarrow & C_2(X', Y; \mathbb{Z}[H]) & \rightarrow 0 \\
& & & \hat{B}_1, \dots, \hat{B}_n & & \hat{B}_1, \dots, \hat{B}_n & \\
& & & \hat{W}_1, \dots, \hat{W}_l & & \hat{W}_1, \dots, \hat{W}_l & \\
& & & \downarrow & & \downarrow & \\
0 & \rightarrow & C_1(Y; \mathbb{Z}[H]) & \rightarrow & C_1(X'; \mathbb{Z}[H]) & \rightarrow & C_1(X', Y; \mathbb{Z}[H]) \rightarrow 0 \\
& & \hat{C}_1^*, \dots, \hat{C}_l^* & \rightarrow & \hat{A}_1^*, \dots, \hat{A}_m^* & & \hat{A}_1^*, \dots, \hat{A}_m^* \\
& & & & \hat{C}_1^*, \dots, \hat{C}_l^* & & \\
& & & \downarrow & & \downarrow & \\
0 & \rightarrow & C_0(Y; \mathbb{Z}[H]) & \rightarrow & C_0(X'; \mathbb{Z}[H]) & \rightarrow & 0. \\
& & \hat{p} & & \hat{p} & &
\end{array}$$

Fox calculus tells us that the boundary map $C_2(X'; \mathbb{Z}[H]) \rightarrow C_1(X'; \mathbb{Z}[H])$ is given by

$$\begin{pmatrix} \varphi \left(\frac{\partial b_j}{\partial a_i} \right) & \varphi \left(\frac{\partial e_*([c_j^*])}{\partial a_i} \right) \\ 0 & * \end{pmatrix}.$$

Thus A is the matrix of the boundary map $C_2(X', Y; \mathbb{Z}[H]) \rightarrow C_1(X', Y; \mathbb{Z}[H])$, and the claim follows. $\square$

6. ALGEBRAIC PROPERTIES OF THE TORSION

In this section, we collect some algebraic properties of the torsion function and their consequences. We begin by describing some known examples which appear as special cases of the torsion for sutured manifolds. We then turn our attention to the “evaluation homomorphism” $H_1(M) \rightarrow H_1(M, R_-(\gamma))$ and prove Proposition 5 from the introduction. Finally, we discuss sutured L-spaces and give the proof of Corollary 6.

6.1. Special Cases of the Torsion. In this section, we summarize some useful special cases of the torsion. These are all “decategorifications” of known facts about sutured Floer homology, although in many cases they admit more elementary proofs as well.

Lemma 6.1. *Let (M, γ) be a product sutured manifold. Denote by $\mathfrak{s}_0$ the canonical vertical Spin^c -structure of (M, γ) and take ω to be the positive orientation on $H_*(M, R_-(\gamma); \mathbb{R}) = 0$. Then*

$$T_{(M, \gamma, \omega)}(\mathfrak{s}) = \begin{cases} 1, & \text{if } \mathfrak{s} = \mathfrak{s}_0, \\ 0, & \text{if } \mathfrak{s} \neq \mathfrak{s}_0. \end{cases}$$

This decategorifies the fact that $SFH(M, \gamma, \mathfrak{s})$ is isomorphic to $\mathbb{Z}$ if $\mathfrak{s} = \mathfrak{s}_0$ and is trivial otherwise [Ju06].

Proof. Let Z be the handle decomposition of (M, γ) with underlying sutured handle complex $(A, S \times I)$ such that $S = R_-(\gamma)$ and $A = S \times I$. Then $s_Z^{-1}(\mathfrak{s}_0) \in \text{Eul}(Z)$ can be represented by the 1-cycle $\theta = \emptyset$. Since there are no handles, we have a canonical identification between $\text{Eul}(Z)$ and $H_1(M)$, and by the definitions of subsection 3.2 in this case $\tau(Z, h, \omega) = h$ for every $h \in \text{Eul}(Z) \cong H_1(M)$. The result follows. $\square$

Lemma 6.2. *Let Y be a closed 3-manifold and let $Y(1)$ be the balanced sutured manifold defined in Example 2.2. Then for every $\mathfrak{s} \in \text{Spin}^c(Y(1))$ we have*

$$T_{Y(1)}(\mathfrak{s}) = \begin{cases} 1 & \text{if } b_1(Y) = 0, \\ 0 & \text{if } b_1(Y) > 0. \end{cases}$$

Proof. Since $R_-(\gamma) = D^2$, the map $p_*: H_1(M) \rightarrow H_1(M, R_-(\gamma))$ is an isomorphism. So the result follows immediately from Proposition 5. $\square$

This also follows from the isomorphism $SFH(Y(1)) \cong \widehat{HF}(Y)$ from [Ju06], together with the corresponding calculation of $\chi(\widehat{HF}(Y, \mathfrak{s}))$ in [OS04a].

Let $L \subset S^3$ be an ordered oriented k -component link. Let $S^3(L)$ be the corresponding balanced sutured manifold as defined in Example 2.3. Then $H = H_1(S^3 \setminus N(L))$ is the free abelian multiplicative group generated by $t_1, \dots, t_k$, where t_k is represented by the meridian of the k -th component of L .

Lemma 6.3. *If $L \subset S^3$ is a k -component link, then given $\mathfrak{s} \in \text{Spin}^c(S^3(L))$ we have*

$$\tau(S^3(L)) = \sum_{h \in H} T_{S^3(L)}(h + \mathfrak{s}) \cdot h \sim \begin{cases} \Delta_L(t_1) & \text{if } k = 1 \\ \Delta_L(t_1, \dots, t_k) \cdot \prod_{i=1}^k (t_i - 1) & \text{if } k \geq 2. \end{cases}$$

Proof. Let $Q = \mathbb{Q}(t_1, \dots, t_k)$. Note that we have a short exact sequence of chain complexes

$$0 \rightarrow C_*(R_-(\gamma); Q) \rightarrow C_*(S^3 \setminus N(L); Q) \rightarrow C_*(S^3 \setminus N(L), R_-(\gamma); Q) \rightarrow 0.$$

From the multiplicativity of torsion (cf. [Tu01, Theorem 1.5]) it follows that

$$\tau(S^3 \setminus N(L), R_-(\gamma); Q) = \tau(S^3 \setminus N(L); Q) \cdot \tau(R_-(\gamma); Q)^{-1}.$$

For $i = 1, \dots, k$ we denote by R_i the component of $R_-(\gamma)$ corresponding to the i -th component of the link L . It follows easily from the definition of the torsion that $\tau(R_i; Q) = (t_i - 1)^{-1}$. We therefore deduce that

$$\begin{aligned} \tau(S^3 \setminus N(L), R_-(\gamma); Q) &= \tau(S^3 \setminus N(L); Q) \cdot \tau(R_-(\gamma); Q)^{-1} = \\ &= \tau(S^3 \setminus N(L); Q) \cdot \prod_{i=1}^k \tau(R_i; Q)^{-1} = \tau(S^3 \setminus N(L); Q) \cdot \prod_{i=1}^k (t_i - 1). \end{aligned}$$

The lemma now follows from the following relation between the torsion and the Alexander polynomial (cf. [Tu01, Theorem 11.8]):

$$\tau(S^3 \setminus N(L); Q) = \begin{cases} \Delta_L(t_1)(t_1 - 1)^{-1}, & \text{if } k = 1 \\ \Delta_L(t_1, \dots, t_k), & \text{if } k \geq 2. \end{cases}$$

□

The following lemma can be viewed as the decategorification of [Ju08, Theorem 1.5].

Lemma 6.4. *Let $K \subset S^3$ be a knot and let R be a genus minimizing Seifert surface for K . Then*

$$\sum_{\mathfrak{s} \in \text{Spin}^c(S^3(R))} T_Y(R)(\mathfrak{s})$$

is equal to the coefficient of $t^{g(R)}$ in the symmetrized Alexander polynomial of K .

Proof. Recall that there exists a Mayer–Vietoris sequence

$$H_1(R) \otimes \mathbb{Z}[t^{\pm 1}] \xrightarrow{A_- - tA_+} H_1(S^3 \setminus N(R)) \otimes \mathbb{Z}[t^{\pm 1}] \rightarrow H_1(S^3 \setminus N(K); \mathbb{Z}[t^{\pm 1}]) \rightarrow 0.$$

Here we endow $H_1(R)$ with any basis and $H_1(S^3 \setminus N(R))$ with the dual basis. Then the symmetrized Alexander polynomial of K equals

$$p(t) = \det(A_- - tA_+) t^{-g(R)}.$$

The coefficient of $t^{g(R)}$ in $p(t)$ is

$$\det(A_+) = |H_1(S^3 \setminus N(R), R_-(\gamma))| = |H_1(S^3(R), R_-(\gamma))|,$$

where we write $|G| = 0$ if G is infinite. With this convention $\epsilon(I_G) = |G|$ for any group G , where $\epsilon : \mathbb{Z}[G] \rightarrow \mathbb{Z}$ is the augmentation map. The lemma now follows immediately from Lemma 6.3 and Proposition 5. □

6.2. Evaluation. In this section, we study the behavior of the torsion under the natural map $p : H_1(M) \rightarrow H_1(M, R_-(\gamma))$. We prove the following statement, which is clearly equivalent to Proposition 5 of the introduction.

Proposition 6.5. *If (M, γ) is a balanced sutured manifold, then*

$$p_*(\tau(M, \gamma)) = \pm I_{H_1(M, R_-(\gamma))},$$

where $I_G \in \mathbb{Z}[G]$ is the sum of all elements in G if G is finite, and is 0 otherwise.

Equivalently, if $K = PD[\ker p_*] \subset H^2(M, \partial M)$, then there is an $\varepsilon \in \{-1, 1\}$ such that for every $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$

$$\sum_{h \in K} T_{(M, \gamma)}(\mathfrak{s} + h) = \begin{cases} \varepsilon & \text{if } H_1(M, R_-(\gamma)) \text{ is finite,} \\ 0 & \text{otherwise.} \end{cases}$$

This result generalizes the fact that $\Delta_K(1) = \pm 1$ whenever $K \subset Y$ is a knot in a homology sphere. Indeed, if we take (M, γ) to be the manifold $Y(K)$ from Example 2.3, then $H_1(M, R_-(\gamma)) = 0$, and p_* induces the map $\mathbb{Z}[t^{\pm 1}] \rightarrow \mathbb{Z}$ which is evaluation at $t = 1$. We also refer to [Tu02, Corollary II 5.2.1], where it is shown that for a null-homologous knot K in a rational homology sphere Y we have $\Delta_K(1) = \pm |H_1(Y)|$.

Proof. By adding product 1-handles, we reduce to the case where $R_-(\gamma)$ is connected. Indeed, if (M', γ') is obtained from (M, γ) by adding a product 1-handle, then $H_1(M', R_-(\gamma')) \cong H_1(M, R_-(\gamma))$, and the diagram

$$\begin{array}{ccc} H_1(M) & \xrightarrow{p_*} & H_1(M, R_-(\gamma)) \\ i_* \downarrow & & \downarrow \cong \\ H_1(M') & \xrightarrow{p'_*} & H_1(M', R_-(\gamma')) \end{array}$$

commutes. Applying Lemma 3.13, we see that if we know that $p'_*(\tau(M', \gamma')) = I_{H_1(M', R_-(\gamma'))}$, then $p_*(\tau(M, \gamma)) = I_{H_1(M, R_-(\gamma))}$ as well.

From now on, we assume that $R_- = R_-(\gamma)$ is connected. Denote the group $H_1(M, R_-)$ by G . Let $\psi : \pi_1(M) \rightarrow G$ be the composition of p_* with the abelianization map, and consider the connected covering map $\pi : \widetilde{M} \rightarrow M$ corresponding to the kernel of ψ . We write $\widetilde{R}_- = \pi^{-1}(R_-)$. Let $\overline{\tau}$ be the maximal abelian torsion of $C_*(\widetilde{M}, \widetilde{R}_-)$, viewed as a module over $\mathbb{Z}[G]$. Then

$$C_*(\widetilde{M}, \widetilde{R}_-) \cong C_*(\hat{M}, \hat{R}_-) \otimes_{\mathbb{Z}[H_1(M)]} \mathbb{Z}[G].$$

It now follows immediately from the proof of Lemma 3.5 that $\overline{\tau} = p_*(\tau(M, \gamma))$. Thus it suffices to show that $\overline{\tau} = I_G$.

As in subsection 3.3, let $T \subset G$ be the torsion subgroup, and pick a splitting $G = F \times T$, where F is a free abelian group. Under the isomorphism

$$\mathbb{Q}[T] \cong \bigoplus_i \mathbb{F}_{\chi_i}$$

I_T maps to the element whose $\mathbb{F}_{\chi_i}$ component is 0 for all non-trivial χ_i and whose component in $\mathbb{F}_{\chi_{\text{id}}} \cong \mathbb{Q}$ is $|T|$. (Here χ_{id} denotes the trivial character.) To show that $\bar{\tau}$ is a multiple of I_G , it suffices to show that the torsion $\tau^{\varphi_i}(M, R_-)$ vanishes whenever either the group F or the character χ_i is nontrivial. Equivalently, we must prove that the complex $C_*(\widetilde{M}, \widetilde{R}_-) \otimes Q(\mathbb{F}_{\chi_i}[F])$ has nontrivial homology.

To this end, we consider the groups $H_0(\widetilde{M})$ and $H_0(\widetilde{R}_-)$. By construction $H_0(\widetilde{M}) = \mathbb{Z}$. Applying the universal coefficient theorem, we see that $H_0(M, Q(\mathbb{F}_{\chi_i}[F])) = 0$ unless $F = 0$ and χ_i is the trivial character. On the other hand, if $e_* : \pi_1(R_-) \rightarrow \pi_1(M)$ is the map induced by the embedding e , then $\psi \circ e_*$ factors as a composition

$$\pi_1(R_-) \rightarrow H_1(R_-) \rightarrow H_1(M) \rightarrow H_1(M, R_-)$$

and is therefore the zero map. It follows that $\pi : \widetilde{R}_- \rightarrow R_-$ is a trivial covering. We assumed R_- is connected, so p_* is a surjection and the deck group of $\widetilde{M}$ is isomorphic to G . Thus $H_0(\widetilde{R}_-) \cong \mathbb{Z}[G]$, from which it follows that $H_0(R_-, Q(\mathbb{F}_{\chi_i}[F])) \cong Q(\mathbb{F}_{\chi_i}[F])$ for any character χ_i .

We now consider the long exact sequence of the pair $(\widetilde{M}, \widetilde{R}_-)$:

$$\rightarrow H_1(M, R_-; Q(\mathbb{F}_{\chi_i}[F])) \rightarrow H_0(R_-; Q(\mathbb{F}_{\chi_i}[F])) \rightarrow H_0(M; Q(\mathbb{F}_{\chi_i}[F])) \rightarrow$$

The middle group in this sequence has rank 1, but the last group is trivial unless $F = 0$ and $\chi_i = 1$. It follows that $H_1(M, R_-; Q(\mathbb{F}_{\chi_i}[F]))$ is nontrivial unless $F = 0$ and χ_i is trivial.

To finish the proof, we need only compute the torsion $\tau^{\varphi_{\text{id}}}(M, R_-)$ when $F = 0$ and φ_{id} is the homomorphism induced by the trivial character. But in this case $C_*(\widetilde{M}) \otimes Q(\mathbb{F}_{\chi_{\text{id}}}[F])$ reduces to the ordinary chain complex $C_*(M, R_-; \mathbb{Q})$. This complex is trivial for $i \neq 1, 2$, so the torsion is $\det(d)$, where $d : C_2(M, R_-) \rightarrow C_1(M, R_-)$ is the boundary map. In other words, $\tau^{\varphi_{\text{id}}}(M, R_-) = \pm |H_1(M, R_-)| = \pm |G|$ as desired. $\square$

6.3. Sutured L -spaces. From the introduction, we recall the following definition.

Definition. We say that (M, γ) is a *sutured L -space* if $SFH(M, \gamma)$ is torsion free and supported in a single $\mathbb{Z}/2$ homological grading.

For sutured L -spaces, the sutured Floer homology is determined by its Euler characteristic; i.e., by the torsion function. In fact, the sutured Floer homology of a sutured L -space has an especially simple form.

Corollary 6.6. *If (M, γ) is a sutured L -space, then for each $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ the group $SFH(M, \gamma, \mathfrak{s})$ is either trivial or isomorphic to $\mathbb{Z}$.*

Proof. Since SFH is supported in a single homological grading, $T_{(M,\gamma)}(\mathfrak{s})$ is equal to the rank of $SFH(M, \gamma, \mathfrak{s})$. Then Proposition 5 shows that

$$\sum_{h \in K} \text{rank } SFH(M, \gamma, \mathfrak{s} + h) \leq 1.$$

This clearly implies the statement above. $\square$

Lemma 6.7. *Suppose that (M, γ) is a balanced sutured manifold and that $S \subset M$ is a nice decomposing surface. Let (M', γ') be the result of decomposing (M, γ) along S . Let O_S be the set of outer Spin^c structures for S , as defined in [Ju08]. If*

$$\bigoplus_{s \in O_S} SFH(M, \gamma, \mathfrak{s})$$

is torsion free and supported in a single $\mathbb{Z}/2$ grading, then (M', γ') is a sutured L -space.

Proof. By the decomposition formula of [Ju08],

$$SFH(M', \gamma') \cong \bigoplus_{s \in O_S} SFH(M, \gamma, \mathfrak{s}).$$

The proof there also gives that $SFH(M', \gamma')$ is supported in a single $\mathbb{Z}/2$ grading. $\square$

Corollary 6.8. *If (M, γ) is a sutured L -space and (M', γ') is obtained by decomposing (M, γ) along a nice surface, then (M', γ') is also a sutured L -space.*

Corollary 6.9. *Suppose that R is a minimal genus Seifert surface of an oriented k -component link $L \subset S^3$. If $\widehat{HFK}(L, g(R) + k - 1)$ is torsion free and supported in one $\mathbb{Z}/2$ grading, then $S^3(R)$ is a sutured L -space.*

Remark. Here we require from any Seifert surface that it intersects a meridian of each component of L geometrically once.

Proof. The sutured manifold $S^3(R)$ is obtained by decomposing $S^3(L)$ along R . By [Ju06], there is an isomorphism $SFH(S^3(L)) \cong \widehat{HFK}(L)$. As shown in [Ju08], the part corresponding to the outer Spin^c structures with respect to R is exactly $\widehat{HFK}(L, g(R) + k - 1)$. $\square$

Corollary 6.10. *If $L \subset S^3$ is a non-split alternating link and R is a minimal genus Seifert surface of L , then $S^3(R)$ is a sutured L -space.*

Proof. The main theorem of [OS03] implies that $\widehat{HFK}(L, g(R) + k - 1)$ is torsion free and supported in a single homological grading, so the result follows from Corollary 6.9. $\square$

We remark that there are many non-alternating links which also satisfy the hypothesis of Corollary 6.9. For example, it is satisfied by all knots of ten or fewer crossings.

7. THE THURSTON NORM FOR SUTURED MANIFOLDS

Let (M, γ) be a sutured manifold. In [Sc89], Scharlemann introduced a natural seminorm on $H_2(M, \partial M; \mathbb{R})$ which generalizes the usual Thurston norm of [Th86]. In this section, we investigate the relation between this norm and the sutured Floer homology.

Definition 7.1. Let (M, γ) be a sutured manifold. Given a properly embedded, compact, connected surface $S \subset M$ let

$$x^s(S) = \max\{0, \chi(S \cap R_-(\gamma)) - \chi(S)\},$$

and extend this definition to disconnected surfaces by taking the sum over the components. Note that $S \cap R_-(\gamma)$ is necessarily a one-dimensional manifold and $\chi(S \cap R_-(\gamma))$ equals the number of components of $S \cap R_-(\gamma)$ which are not closed. Equivalently, we have

$$x^s(S) = \max \left\{ 0, -\chi(S) + \frac{1}{2}|S \cap s(\gamma)| \right\}.$$

For $\alpha \in H_2(M, \partial M)$ let

$$x^s(\alpha) = \min\{x^s(S) : S \subset M \text{ is properly embedded and } [S, \partial S] = \alpha\}.$$

Theorem 7.2. [Sc89] *Let (M, γ) be a sutured manifold. Then the function*

$$x^s : H_2(M, \partial M) \rightarrow \mathbb{Z}_{\geq 0}$$

defined above has the following two properties:

- (1) $x^s(n\alpha) = |n| \cdot x^s(\alpha)$ for all $n \in \mathbb{Z}$ and $\alpha \in H_2(M, \partial M)$,
- (2) $x^s(\alpha + \beta) \leq x^s(\alpha) + x^s(\beta)$ for all $\alpha, \beta \in H_2(M, \partial M)$.

It follows that x^s extends to a continuous map $x^s : H_2(M, \partial M; \mathbb{R}) \rightarrow \mathbb{R}_{\geq 0}$ which is convex and linear on rays from the origin. Put differently, x^s is a seminorm on $H_2(M, \partial M; \mathbb{R})$. It is called the *sutured Thurston norm*.

Example. Let Y be a closed 3-manifold and let $(M, \gamma) = Y(1)$ be Y with an open ball removed and having a connected suture. Then we can identify $H_2(Y; \mathbb{R})$ with $H_2(Y(1), \partial Y(1); \mathbb{R})$. It is straightforward to see that under this identification the Thurston norms of Y and $Y(1)$ agree.

Example. Let $K \subset S^3$ a knot and let $(M, \gamma) = S^3(K)$ be the associated sutured manifold with two meridional sutures. If $\alpha \in H_2(M, \partial M)$ is a generator, then $x^s(\alpha) = 2g(K)$. Note that this differs from the usual Thurston norm x of M , which satisfies $x(\alpha) = 2g(K) - 1$ for a non-trivial knot.

The previous example is in fact a special case of the following elementary lemma.

Lemma 7.3. *Let Y be a closed 3-manifold and let $L \subset Y$ be an l -component link with meridians $\mu_1, \dots, \mu_l$. Suppose that L has no trivial components. Let $Y(L) = (M, \gamma)$ be the corresponding sutured manifold. Given $h \in H_2(M, \partial M)$ we have*

$$x^s(h) = x(h) + \sum_{i=1}^l |\langle h, \mu_i \rangle|,$$

where $x(h)$ denotes the ordinary Thurston norm on $H_2(M, \partial M)$.

The following proposition should be compared with [Ju08b, Theorem 5.1] and [Ju08b, Proposition 7.5]. For all the necessary definitions also see [Ju08b].

Proposition 7.4. *Suppose that the taut balanced sutured manifold (M, γ) is reduced, horizontally prime, and $H_2(M) = 0$. Then x^s is a norm.*

Proof. Assume there exists an $\alpha \neq 0$ in $H_2(M, \partial M)$ with $x^s(\alpha) = 0$. This implies that there exists a connected homologically non-trivial orientable surface $(S, \partial S) \subset (M, \partial M)$ with $x^s(S) = 0$. Hence

$$-\chi(S) \leq -\chi(S) + \chi(S \cap R_-(\gamma)) \leq 0.$$

So $\chi(S) \geq 0$ and S is either S^2, T^2, D^2 , or $S^1 \times I$. Since $[S, \partial S] \neq 0$ and $H_2(M) = 0$ the surface S is not S^2 or T^2 . Furthermore, we can assume that $S \cap \gamma$ consists of arcs connecting $R_-(\gamma)$ and $R_+(\gamma)$.

Now suppose that $S = D^2$. Then $\chi(S \cap R_-(\gamma))$ is 0 or 1. In the latter case S would be a homologically non-trivial product disc, contradicting the assumption that (M, γ) is reduced. In the former case S is a compressing disk for $R(\gamma)$. Since (M, γ) is taut $R(\gamma)$ is incompressible, so ∂S bounds a disk S' in $R(\gamma)$. Now $S \cup (-S')$ is a sphere, which has to bound a D^3 since M is irreducible. But then $[S, \partial S] = 0$, a contradiction.

Finally, assume that S is an annulus. Then $\chi(S \cap R_-(\gamma)) = 0$. Since (M, γ) is reduced we know that S can not be a product annulus. So suppose that $\partial S \subset R$, where R is either $R_-(\gamma)$ or $R_+(\gamma)$. Pick a product neighborhood $S \times [0, 1]$ and let

$$R' = R \setminus (\partial S \times (0, 1)) \cup (S \times 0) \cup (S \times 1).$$

Then R' is homologous to R , $\partial R' = \partial R$ and $\chi(R') = \chi(R)$. Hence R' is a horizontal surface. Note that R' is not parallel to R . If R' were parallel to $R(\gamma) \setminus R$ then $\partial S \times [0, 1]$ would give rise to a non-trivial product annulus. So the existence of R' would contradict our assumption that (M, γ) is horizontally prime. $\square$

Definition 7.5. Let $S(M, \gamma) = \{\mathfrak{s} \in \text{Spin}^c(M, \gamma) : SFH(M, \gamma, \mathfrak{s}) \neq 0\}$ be the support of $SFH(M, \gamma)$. If $\alpha \in H_2(M, \partial M; \mathbb{R})$, we define

$$z(\alpha) = \max\{\langle \mathfrak{s} - \mathfrak{t}, \alpha \rangle : \mathfrak{s}, \mathfrak{t} \in S(M, \gamma)\}.$$

Remark. In [Ju08b, Section 8] another “seminorm” y on $H_2(M, \partial M; \mathbb{R})$ was constructed using sutured Floer homology. The function y satisfies all properties of a seminorm except that $y(\alpha) \neq y(-\alpha)$ can happen. It is straightforward to see that

$$z(\alpha) = \frac{1}{2}(y(\alpha) + y(-\alpha)).$$

The following proposition proves the second statement of Theorem 4.

Proposition 7.6. *Let (M, γ) be an irreducible balanced sutured manifold such that all boundary components of M are tori. Then $z = x^s$.*

Proof. First note that by standard arguments it suffices to show the equality of norms for integral classes.

Now assume that each component of ∂M has exactly two sutures. Observe that we can assign to (M, γ) a link L in a 3-manifold Y which is obtained by Dehn filling ∂M such that the μ_i become meridians of the filling tori, see [Ju06, Example 2.4]. Then by [Ju08, Remark 8.5] we can assign to each $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ a relative first Chern class $c_1(\mathfrak{s}) \in H^2(M, \partial M)$ in such a way that the set $\{c_1(\mathfrak{s}) : \mathfrak{s} \in S(M, \gamma)\}$ is symmetric about the origin. Then by [Ju08, Remark 8.5] for every $h \in H_2(M, \partial M)$

$$\max\{\langle c_1(\mathfrak{s}), h \rangle : \mathfrak{s} \in S(M, \gamma)\} = x(h) + \sum_{i=1}^l |\langle h, \mu_i \rangle|.$$

Since the image of $S(M, \gamma)$ is centrally symmetric, this is equivalent to saying

$$\max\{\langle \mathfrak{s} - \mathfrak{t}, h \rangle : \mathfrak{s}, \mathfrak{t} \in S(M, \gamma)\} = x(h) + \sum_{i=1}^l |\langle h, \mu_i \rangle|.$$

Note that the left hand side is just $z(h)$. If L is not the unknot, then L does not have trivial components since M is irreducible. So by Lemma 7.3 the right hand side is exactly $x^s(h)$. The proposition is obviously true if L is the unknot.

Now consider the general case that each component of ∂M has at least two sutures. We can reduce this case to the case treated above using Proposition [Ju08b, Proposition 9.2]. $\square$

Remark. Link Floer homology of a link $L \subset S^3$ was defined in [OS08a]. It agrees with the sutured Floer homology of the sutured manifold $S^3(L)$ introduced in Example 2.3. In [OS08b] it is shown that if L has no trivial components then the link Floer homology of L determines the Thurston norm of the link complement. In light of the above theorem it is perhaps a better point of view to observe that twice the link Floer polytope equals the dual of the sutured Thurston polytope of $S^3(L)$, which then determines the ordinary Thurston polytope of $S^3(L)$ via Lemma 7.3.

The following theorem is exactly the first part of Theorem 4.

Theorem 7.7. *Let (M, γ) be an irreducible balanced sutured manifold. Then $z \leq x^s$.*

We will later see in Proposition 7.12 that the inequality of Theorem 7.7 is strict in general.

In order to prove Theorem 7.7 we consider the double of the sutured manifold (M, γ) along $R(\gamma)$. More precisely, the double DM of (M, γ) is obtained from the disjoint union of M and $-M$ by identifying the two copies of $R(\gamma)$ via the identity map. The boundary of DM is a union of tori; each torus is the double of a component of γ . This operation was first used by Gabai [Ga83].

A theorem of Cantwell and Conlon relates the sutured Thurston norm on (M, γ) to the Thurston norm of the double. To be precise, suppose (M, γ) is a sutured manifold, and let $X = DM$ be the double of M along $R(\gamma)$. There is a natural “doubling map” $D_* : H_2(M, \partial M; \mathbb{R}) \rightarrow H_2(X, \partial X; \mathbb{R})$.

Theorem 7.8. [CC06, Theorem 2.3] *We have $x(D_*(\alpha)) = 2x^s(\alpha)$.*

Here x denotes the usual Thurston norm on $H_2(X, \partial X; \mathbb{R})$ and x^s is the sutured Thurston norm on $H_2(M, \partial M; \mathbb{R})$.

Definition 7.9. We make X into a sutured manifold (X, γ_X) in the following canonical way. Let the components of γ be $\gamma_1, \dots, \gamma_l$. For each component γ_i of γ choose two parallel, oppositely oriented arcs m_i and m'_i that connect $R_+(\gamma)$ and $R_-(\gamma)$. Then on the torus $D\gamma_i \subset \partial X$ the sutures are $\mu_i = m_i \cup (-m_i)$ and $\mu'_i = m'_i \cup (-m'_i)$. These sutures are well defined up to isotopy. Let γ_X be a regular neighborhood of $\bigcup_{i=1}^l (\mu_i \cup \mu'_i)$ inside ∂X .

Lemma 7.10. *Let (M, γ) be an irreducible balanced sutured manifold and let (X, γ_X) be its double. Then for all $\alpha \in H_2(M, \partial M; \mathbb{R})$ we have*

$$2x^s(\alpha) = \max\{\langle \mathfrak{s} - \mathfrak{t}, D_*(\alpha) \rangle : \mathfrak{s}, \mathfrak{t} \in S(X, \gamma_X)\}.$$

Proof. First note that by standard arguments it suffices to show the equality when α is an integral class.

Since X is irreducible and has only toroidal boundary components, by Proposition 7.6 we have

$$\max\{\langle \mathfrak{s} - \mathfrak{t}, h \rangle : \mathfrak{s}, \mathfrak{t} \in S(X, \gamma_X)\} = x^s(h)$$

for any $h \in H_2(X, \partial X)$.

We claim that if $h = D_*(\alpha)$ for some $\alpha \in H_2(M, \partial M)$ then $\langle h, \mu_i \rangle = 0$ for every $1 \leq i \leq l$. To see this choose a surface S representing α . We may assume that $S \cap \gamma_i$ consists of a collection of parallel arcs. If we take m_i and m'_i parallel to these arcs, then $\partial S \cap \mu_i$ and $\partial S \cap \mu'_i$ are empty. Combining this with the fact that $x(h) = 2x^s(\alpha)$, we obtain the statement of the lemma. $\square$

The surface $R(\gamma)$ defines an oriented surface $R \subset X$. Note that R has the orientation coming from $R(\gamma)$, *not* the induced orientation coming from ∂M . In particular, the homology class represented by R is twice the class of $R_-(\gamma)$. It is easy to see that R is a nice decomposing surface for (X, γ_X) in the sense of [Ju08b]. Let $O_R \subset \text{Spin}^c(X, \gamma_X)$ be the set of outer Spin^c structures for R .

Lemma 7.11. *Let (M, γ) be a taut balanced sutured manifold. Then for all $\alpha \in H_2(M, \partial M; \mathbb{R})$ we have*

$$2z(\alpha) = \max \{ \langle \mathfrak{s} - \mathfrak{t}, D_*(\alpha) \rangle : \mathfrak{s}, \mathfrak{t} \in O_R \cap S(X, \gamma_X) \}.$$

Proof. First note that by standard arguments it suffices to show the equality when α is an integral class.

Now note that if we decompose (X, γ_X) along R we get $(M', \gamma') = (M, \gamma) \sqcup (-M, \gamma)$. $\text{Spin}^c(M', \gamma')$ is naturally identified with $\text{Spin}^c(M, \gamma) \times \text{Spin}^c(-M, \gamma)$. If $i : M' \rightarrow X$ is the inclusion, there is a map $f_R : \text{Spin}^c(M', \gamma') \rightarrow O_R$ with the property that $f_R(\mathfrak{s}) - f_R(\mathfrak{t}) = i_*(\mathfrak{s} - \mathfrak{t})$, as in [Ju08b, Prop. 5.4]. (Here we view $\mathfrak{s} - \mathfrak{t}$ as an element of $H_1(X)$, or equivalently, use $PD \circ i_* \circ PD$ in place of i_* .) The decomposition theorem of [Ju08b] implies that

$$f_R(S(M', \gamma')) = O_R \cap S(X, \gamma_X).$$

Clearly

$$S(M', \gamma') = S(M, \gamma) \times S(-M, \gamma).$$

Recall that there is a bijection $\text{Spin}^c(M, \gamma) \rightarrow \text{Spin}^c(-M, \gamma)$ which sends a nonvanishing vector field v to $-v$. By Proposition 2.12, we have $S(-M, \gamma) = -S(M, \gamma)$. Thus each element of $O_R \cap S(X, \gamma_X)$ can be written as $\mathfrak{s} = f_R(\mathfrak{s}_1, -\mathfrak{s}_2)$, where $\mathfrak{s}_1, \mathfrak{s}_2 \in \text{Spin}^c(M, \gamma)$. So for $\mathfrak{s}, \mathfrak{t} \in O_R \cap S(X, \gamma_X)$ we have

$$\begin{aligned} \langle \mathfrak{s} - \mathfrak{t}, D_*(\alpha) \rangle &= \langle f_R(\mathfrak{s}_1, -\mathfrak{s}_2) - f_R(\mathfrak{t}_1, -\mathfrak{t}_2), D_*(\alpha) \rangle \\ &= \langle i_*(\mathfrak{s}_1 - \mathfrak{t}_1, \mathfrak{t}_2 - \mathfrak{s}_2), D_*(\alpha) \rangle \\ &= \langle \mathfrak{s}_1 - \mathfrak{t}_1, \alpha \rangle + \langle \mathfrak{s}_2 - \mathfrak{t}_2, \alpha \rangle. \end{aligned}$$

In particular, we see that

$$\max \{ \langle \mathfrak{s} - \mathfrak{t}, D_*(\alpha) \rangle : \mathfrak{s}, \mathfrak{t} \in O_R \cap S(X, \gamma_X) \} = 2 \cdot \max \{ \langle \mathfrak{s} - \mathfrak{t}, \alpha \rangle : \mathfrak{s}, \mathfrak{t} \in S(M, \gamma) \}.$$

The right hand side is by definition $2z(\alpha)$. $\square$

We are now finally ready to complete the proof of Theorem 7.7.

Proof of Theorem 7.7. If (M, γ) is taut, this is an immediate consequence of Lemmas 7.10 and 7.11. Now suppose that (M, γ) is not taut. Since M is irreducible [Ju06, Proposition 9.18] implies that $SFH(M, \gamma) = 0$. So $z = 0$, and the inequality is obviously true. $\square$

Remark. We introduce the notation B_{x^s} for the unit norm ball of x^s and B_z for the unit norm ball of z . Let

$$P = \{ c_1(\mathfrak{s}) : \mathfrak{s} \in S(X, \gamma_X) \} \subset H^2(X, \partial X; \mathbb{R})$$

be the sutured Floer polytope of (X, γ_X) , and for any $h \in H_2(X, \partial X)$ let P_h be the outer face of P in the direction h ; i.e.

$$P_h = \{ x \in P : \langle x, h \rangle \text{ is maximal} \}.$$

Finally, let $Q \subset H^2(M, \partial M; \mathbb{R})$ be the sutured Floer polytope of (M, γ) , see [Ju08b]. Then Lemmas 7.10 and 7.11 tell us that

$$B_{2x^s} = P^* \cap \text{Im}(D_*)$$

and

$$\begin{aligned} B_{2z} &= \left(\frac{P_\rho + P_{-\rho}}{2} \right)^* \cap \text{Im}(D_*) \\ &= (Q + (-Q))^*, \end{aligned}$$

where $\rho = [R]$ and where $P_\rho + P_{-\rho}$ denotes the Minkowski sum of P_ρ and $P_{-\rho}$. Of course $B_{2x^s} = B_{x^s}/2$ and $B_{2z} = B_z/2$.

When (M, γ) is a rational homology product, the relation between these two sets can be described as follows. Let $i : R_-(\gamma) \rightarrow X$ be the inclusion, and let $A \subset H_1(X; \mathbb{R})$ be the image of i_* . Let $C \subset H_1(X; \mathbb{R})$ be the one dimensional space spanned by the double of an arc joining $R_-(\gamma)$ to $R_+(\gamma)$. Then $H_1(X; \mathbb{R}) = A \oplus C$.

There is a natural “reflection” $r : X \rightarrow X$ which acts as the identity on A and by multiplication by -1 on C . The image of D_* is fixed by r_* . It follows that C is orthogonal to $\text{Im } D_*$; i.e., the pairing $H_1(X; \mathbb{R}) \times H_2(X, \partial X; \mathbb{R}) \rightarrow \mathbb{R}$ restricted to $C \times \text{Im } D_*$ vanishes. The action of r_* exchanges the faces P_ρ and $P_{-\rho}$. Composing the central symmetry of P with r_* gives an involution on P_ρ . (This realizes the central symmetry of B_z .) It follows that $P_{-\rho}$ is a translate of P_ρ in the direction of C .

Since C is orthogonal to $\text{Im}(D_*)$, restricting our attention to the image of D_* has the effect of projecting P in the direction of C . The norms x^s and z will agree if and only if the image of P under this projection is equal to P_ρ . Somewhat surprisingly, this is not always the case.

Proposition 7.12. *For the sutured manifold (M, γ) given in [CC06, Example 2] we have $x^s \neq z$.*

Proof. In [CC06] the unit ball B_{x^s} is computed explicitly. To get B_z one proceeds by computing $T_{(M, \gamma)}$, which is straightforward since M is a genus three handlebody. It is also easy to see that (M, γ) is a rational homology product. Now to obtain the sutured Floer homology polytope

$$P(M, \gamma, t) = \text{conv}\{c_1(\mathfrak{s}, t) : \mathfrak{s} \in S(M, \gamma)\} \subset H^2(M, \partial M; \mathbb{R})$$

one uses the adjunction inequality of [Ju08b] for the decomposing discs given in [CC06], each of which intersects $s(\gamma)$ in four points. Note that for $\mathfrak{s}, \mathfrak{t} \in \text{Spin}^c(M, \gamma)$ we have $c_1(\mathfrak{s}, t) - c_1(\mathfrak{t}, t) = 2(\mathfrak{s} - \mathfrak{t})$. Hence after symmetrizing one can just observe that $B_z \neq B_{x^s}$. $\square$

Remark. In the above example the polytope P is four-dimensional and is composed of three layers in the ρ direction. The opposite faces P_ρ and $P_{-\rho}$ are in the two outer layers, and they are both smaller than the middle layer, which is exactly $B_{2x^s}^*$. This

picture shows that this is in some sense the smallest possible counterexample where M is a handlebody, since on a genus two handlebody we would have $P = P_\rho \cup P_{-\rho}$.

Let (M, γ) be a balanced sutured manifold such that $H_2(M) = 0$. In [Ju08, Proposition 8.10] the second author showed that if $(M, \gamma) \rightsquigarrow^S (M', \gamma')$ is a decomposition along a “nice” product annulus S , then $SFH(M, \gamma) \cong SFH(M', \gamma')$. Theorem 7.7 permits us to extend this result to irreducible balanced sutured manifolds with arbitrary second homology.

Proposition 7.13. *Suppose that (M, γ) is an irreducible balanced sutured manifold. Let $S \subset (M, \gamma)$ be a product annulus such that at least one component of ∂S is non-zero in $H_1(R(\gamma))$, or both components of ∂S are boundary-coherent in $R(\gamma)$. If S gives a surface decomposition $(M, \gamma) \rightsquigarrow^S (M', \gamma')$, then*

$$SFH(M, \gamma) \cong SFH(M', \gamma').$$

Proof. In both cases we can orient S such that ∂S is boundary-coherent in $R(\gamma)$. Note that (M', γ') is taut if and only if (M, γ) is, and both are irreducible. So if (M', γ') is not taut, then by [Ju06, Proposition 9.18] we have

$$SFH(M, \gamma) = 0 = SFH(M', \gamma').$$

Now assume that (M', γ') is taut, then $SFH(M', \gamma') \neq 0$ by [Ju08, Theorem 1.4]. [Ju08, Theorem 1.3] implies that

$$SFH(M', \gamma') \cong \bigoplus_{\mathfrak{s} \in O_S} SFH(M, \gamma, \mathfrak{s}).$$

So $O_S \cap S(M, \gamma) \neq \emptyset$. Let $\mathfrak{s}_0 \in O_S \cap S(M, \gamma)$, then for any $\mathfrak{s} \in \text{Spin}^c(M, \gamma)$ we have $\langle \mathfrak{s} - \mathfrak{s}_0, [S] \rangle = 0$ if and only if $\mathfrak{s} \in O_S$. Obviously, $x^s([S]) = 0$, so by Theorem 7.7 $z([S]) = 0$. Thus $\langle \mathfrak{s} - \mathfrak{s}_0, [S] \rangle = 0$ for every $\mathfrak{s} \in S(M, \gamma)$. So $S(M, \gamma) \subset O_S$, and hence $SFH(M, \gamma) \cong SFH(M', \gamma')$. $\square$

8. EXAMPLES AND APPLICATIONS

We conclude with some sample computations of the torsion and/or the sutured Floer homology, with emphasis on the case where (M, γ) is the complement of a Seifert surface $R \subset S^3$.

Example 8.1. Suppose that $R \subset S^3$ is an embedded annulus. Then ∂R consists of two parallel copies of a knot K with some linking number n corresponding to the framing of the annulus. The complementary sutured manifold $S^3(R)$ is homeomorphic to $S^3 \setminus N(K)$. Its boundary is a torus with two sutures, each representing the homology class $\ell + nm$ with respect to the canonical basis on $H_1(\partial(S^3 \setminus N(K)))$. Let K_n be the manifold obtained by filling this homology class (*i.e.*, by performing $n/1$ Dehn

surgery on K), and let $K(n) \subset K_n$ be the core circle of the filling. Then $SFH(S^3(R))$ is isomorphic to $\widehat{HFK}(K(n))$, from which it follows that

$$\tau(S^3(R)) = \Delta_K(t) \cdot \frac{t^n - 1}{t - 1}.$$

Note that when $n = 0$, the torsion vanishes, regardless of what K is. The group $\widehat{HFK}(K(n))$ has been studied by Eftekhary [Ef05] (in the case $n = 0$) and by Hedden [He05], who gives a complete calculation in terms of the groups $HFK^-(K)$. In particular, $\widehat{HFK}(K(0))$ is nontrivial unless K is the unknot.

Example 8.2. Suppose M is a solid torus, and that γ consists of $2n$ parallel curves on ∂M , each of which represents p times the generator of $H_1(M)$. $SFH(M, \gamma)$ was computed by the second author in [Ju08b]; its Euler characteristic is given by

$$\tau(M, \gamma) \sim \frac{(t^p - 1)^n}{t - 1}.$$

The homology in each Spin^c structure is a free module of rank equal to the Euler characteristic. An important special case is when $n = 1$. In this case $(M, \gamma) = S^3(R)$, where R is a twisted band with p full twists (in other words, an unknotted annulus in S^3 with framing p .) $SFH(M, \gamma)$ is supported in p consecutive Spin^c structures, each containing a single copy of $\mathbb{Z}$.

Example 8.3. Suppose $K = K(p, q) \subset S^3$ is the two-bridge knot or link corresponding to the fraction p/q . The set of Seifert surfaces for K has been classified up to isotopy by Hatcher and Thurston [HT85]. Any such surface is obtained as a Murasugi sum of twisted bands. By [Ju08, Cor. 8.8] and [Ju08b, Theorem 5.11] the sutured Floer polytope is a rectangular prism, the length of whose sides is determined by the number of twists in the corresponding band. These, in turn, are given by the coefficients of the unique continued fraction expansion of p/q all of whose terms are even [HT85]. For example, the knot $K(56, 15)$ has continued fraction expansion

$$\frac{56}{15} = 4 - \frac{1}{4 - \frac{1}{4}}.$$

Any Seifert surface R of K is a Murasugi sum of three twisted bands, each with two full twists. $SFH(S^3(R)) \cong \mathbb{Z}^8$ is supported at the vertices of a $2 \times 2 \times 2$ cube.

Whenever two of the bands have more than one twist, K will have more than one Seifert surface. The calculation above shows that these surfaces cannot be distinguished by their sutured Floer polytope alone. In contrast, we have the following.

Theorem. [HJS08] *There exist two minimal genus Seifert surfaces R_1 and R_2 for $K(17, 4)$ which can be distinguished by combining sutured Floer homology with the Seifert form. More precisely, there does not exist an orientation-preserving diffeomorphism of the pairs (S^3, R_1) and (S^3, R_2) .*

By Corollary 6.10 the groups $SFH(S^3(R_i), \mathfrak{s}), i = 1, 2$ for $\mathfrak{s} \in \text{Spin}^c(S^3(R_i))$ are determined by $T_{S^3(R_i)}(\mathfrak{s})$ since two-bridge knots are alternating. Hence it is straightforward to modify the proof of the theorem to show that the Seifert surfaces can also be distinguished by using torsion and the Seifert form.




FIGURE 1. The pretzel knot $P(2r+1, 2s+1, 2t+1)$ with $2r+1, 2s+1$, and $2t+1$ half twists

Example 8.4. The pretzel knot $P(2r+1, 2s+1, 2t+1)$ has an obvious Seifert surface R , as shown in Figure 1 (a). A natural pair of compressing disks A and B for the handlebody $M = S^3 \setminus N(R)$ is shown in Figure 1 (b). Cutting M along these disks and using the Seifert–van Kampen theorem give an isomorphism between $\pi_1(M)$ and the free group generated by a and b . If α is a curve on ∂M , we can read off the word it represents in $\pi_1(M)$ by traversing α and recording its intersections with ∂A and ∂B .

Suppose r, s and t are all positive and that R is oriented so that the uppermost region in the figure belongs to $R_-(\gamma)$. Put p in this region, and let α be a loop which runs from p down the left-hand strip, and back up via the middle strip. Similarly, let β be a loop which runs down the right-hand strip and back up the middle, so that $\pi_1(R_-(\gamma), p)$ is generated by α and β . The reader can easily verify that

$$\alpha = a^{r+1}(b^{-1}a)^sb^{-1} \quad \beta = b^{-t}(b^{-1}a)^sb^{-1}.$$

Evaluating the 2×2 determinant

$$\begin{bmatrix} \partial\alpha/\partial a & \partial\beta/\partial a \\ \partial\alpha/\partial b & \partial\beta/\partial b \end{bmatrix}$$

we find that $\tau(S^3(R))$ is supported on a hexagon, as illustrated in Figure 2. With respect to the natural basis given by a and b , the sides of the hexagon have slope $0, 1$ and ∞ . Parallel sides have the same length, and the sides are of length $r+1, t+1$,

and $s + 1$. The coefficient of the torsion at each lattice point in the hexagon is 1, and the sutured Floer homology consists of a single copy of $\mathbb{Z}$ at each lattice point since the pretzel knot is alternating.




FIGURE 2. Support of $\tau(S^3(R))$ for $r, s, t > 0$.

The case where $r, s > 0$ and $t < 0$ can be treated similarly. We distinguish two subcases, depending on whether $|2t - 1|$ is less than $\min(2r + 1, 2s + 1)$, or greater. In the first, the coefficients of $\tau(S^3(R))$ take on both positive and negative signs. The torsion is supported on a “bowtie”, as shown in Figure 3. The coefficient of the torsion is 1 at each lattice point in the rectangle, and -1 at each lattice point in the two triangles. In the second case, the support is a nonconvex hexagon, as illustrated in Figure 4. The coefficient of the torsion is -1 at each lattice point in the hexagon. To determine the sutured Floer homology, we compare with the calculation of $\widehat{HFK}(P(2r + 1, 2s + 1, 2t + 1))$ given in [OS04c]. In both cases, the top group in the knot Floer homology is torsion free and its rank is equal to the number of vertices in the support of $\tau(S^3(R))$. It follows that $SFH(S^3(R))$ has rank one at each vertex in the support and is trivial elsewhere.





FIGURE 3. Support of $\tau(S^3(R))$ for $r, s > 0$ and $-\min(r, s) \leq t \leq 0$.

Example 8.5. The three-component pretzel link $P(2r, 2s, 2t)$ has a Seifert surface R similar to that shown in Figure 1, and $\tau(S^3(R))$ can be computed as in the previous

FIGURE 4. Support of $\tau(S^3(R))$ for $r, s > 0$ and $t \leq -\min(r, s)$.

example. When r, s, t are all positive, the torsion is again supported on a hexagon with sides of slopes 0, 1 and ∞ . However in this case parallel sides of the hexagon do not have the same lengths. Instead, the sides have lengths $r+1, t, s+1, r, t+1, s$ as we go around the hexagon. This gives a simple family of examples for which the torsion does not exhibit any symmetry. The phenomenon is already evident for $P(2, 2, 2)$. In this case, the hexagon degenerates to a triangle supported at three vertices in the plane. With respect to the standard basis a, b , these vertices can be taken to be $(0, 0)$, $(1, 0)$ and $(1, 1)$. Note that for $r, s, t > 0$, $P(2r, 2s, 2t)$ is an alternating link, and hence $SFH(S^3(R))$ has the same support as the torsion. Thus the sutured Floer polytope is asymmetric as well.

Example 8.6. Seifert surfaces of small knots. Let K be a knot in S^3 and suppose R is a Seifert surface for K . Among knots with nine crossings or fewer, most are either two-bridge or fibred. (See *e.g.* the tables in [Ka96] or [CL09].) If K is fibred, $SFH(S^3(R)) \cong \mathbb{Z}$; if it is two-bridge, $SFH(S^3(R))$ was determined in Example 8.3. The remaining knots all have a unique Seifert surface R by [Ka05]. We briefly describe the groups $SFH(S^3(R))$. They fall into two broad classes, as well as a few knots with more interesting homology.

- The knots $9_{16}, 9_{37}$, and 9_{46} have $SFH(S^3(R)) \cong SFH(A_2)$, where A_2 is an unknotted annulus with two full twists.
- The knots $8_{15}, 9_{25}, 9_{39}, 9_{41}$, and 9_{49} have $SFH(S^3(R)) \cong SFH(S^3(R_{2,2,2}))$, where $R_{2,2,2}$ is the Seifert surface of the $(2, 2, 2)$ pretzel link.
- The knot 9_{35} is $P(3, 3, 3)$. Its sutured Floer polytope is a hexagon with sides of length 2.
- The knot 9_{38} is the only knot with fewer than 10 crossings whose sutured Floer polytope is 3-dimensional. The polytope is contained in a $2 \times 2 \times 2$ cube, with $\mathbb{Z}$ summands at five of the vertices: $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 1)$, $(1, 1, 0)$, and $(1, 0, 1)$.

For all of these knots, the top group in knot Floer homology is torsion free and supported in a single homological grading, so SFH is determined by the torsion.

Example 8.7. The four-strand pretzel link $L = P(n, -n, n, -n)$ has a genus one Seifert surface analogous to the one shown in Figure 1. The multivariable Alexander polynomial of this link is 0, but a calculation similar to the one in Example 4 shows that the torsion polytope is a “pinwheel” consisting of four square pyramids, each with side length n . It follows that the rank of $HFK(L)$ in the top Alexander grading is at least

$$4 \sum_{k=1}^n k^2 = \frac{2n(n+1)(2n+1)}{3}.$$




FIGURE 5. Sutured genus two handlebody with no disk decomposition.

Example 8.8. We conclude by using the torsion to give an example of a phenomenon first observed by Goda [Go94]. Namely, there exist sutured manifolds whose total space is a genus two handlebody, but which are not disk-decomposable. Consider the two pairs of pants illustrated in Figure 8. We consider the genus two surface obtained by gluing the two pairs of pants glued along the corresponding boundary curves by identifying the corresponding numbers $1, 2, \dots, 12$. Let M be the handlebody in which the curves labeled A and B bound compressing disks, and let $s(\gamma)$ be the curve shown in the figure. Then we easily compute

$$\tau(M, \gamma) \sim 2a - 3 + 2a^{-1}.$$

Proposition 8.1. (M, γ) is not disk-decomposable.

Proof. Suppose we decompose (M, γ) along a disk D to obtain a sutured manifold (M', γ') . If ∂D is a non-separating curve in ∂M , then M' is homeomorphic to $S^1 \times D^2$. If (M', γ') were taut, then by [Ju08b], $SFH(M', \gamma')$ would be isomorphic to the restriction of $SFH(M, \gamma)$ to those Spin^c structures which are extremal with respect to

evaluation on $[D]$. It follows that either $\tau(M', \gamma') = 0$, $\tau(M', \gamma') \sim 2$, or $\tau(M', \gamma') \sim 2t - 3 + 2t^{-1}$. Comparing with Example 2, we see that none of these are the torsion of a taut sutured manifold whose total space is the solid torus.

Similarly, if ∂D is a separating curve, then M' is homeomorphic to the disjoint union of two solid tori M_1 and M_2 , and

$$SFH(M, \gamma) \cong SFH(M', \gamma') \cong SFH(M_1, \gamma_1) \otimes SFH(M_2, \gamma_2).$$

Again, comparing $\tau(M, \gamma)$ with Example 8.2 shows that this is not possible. $\square$

REFERENCES

- [CC06] J. Cantwell and L. Conlon, *The sutured Thurston norm*, math.GT/060653.
- [Ce70] J. Cerf, *La stratification naturelle des espaces de fonctions différentiables réelles et le théorème de la pseudo-isotopie*, Publ. Math. I.H.É.S. **39** (1970), 5–173.
- [CL09] J.C. Cha and C. Livingston, *KnotInfo: Table of Knot Invariants*, <http://www.indiana.edu/~knotinfo>, 2009.
- [CF77] R. Crowell and R. Fox, *Introduction to knot theory*, reprint of the 1963 original. Graduate Texts in Mathematics, No. 57, Springer-Verlag, New York-Heidelberg, 1977.
- [Ef05] E. Eftekhary, *Longitude Floer homology and the Whitehead double*, Algebr. Geom. Topol. **5** (2005), 1389–1418.
- [Ga83] D. Gabai, *Foliations and the topology of 3-manifolds*, J. Differential Geometry **18** (1983), 445–503.
- [Ga86] D. Gabai, *Detecting fibred links in S^3* , Comment. Math. Helv. **61** (1986), no. 4, 519–555.
- [Go94] H. Goda, *A construction of taut sutured handlebodies which are not disk decomposable*, Kobe J. Math. **11** (1994), no. 1, 107–116.
- [GS08] H. Goda and T. Sakasai, *Homology cylinders in knot theory*, math.GT/0807.4034.
- [HT72] S. Halperin and D. Toledo, *Stiefel-Whitney homology classes*, Ann. of Math **96** (1972), 511–525.
- [HT85] A. Hatcher and W. Thurston, *Incompressible surfaces in 2-bridge knot complements*, Invent. Math. **79** (1985), no. 2, 225–246.
- [He05] M. Hedden, *On Knot Floer Homology and Cabling*, Algebraic and Geometric Topology **5** (2005), 1197–1222.
- [HJS08] M. Hedden, A. Juhász, and S. Sarkar, *On sutured Floer homology and the equivalence of Seifert surfaces*, math.GT/0811.0178.
- [Ho82] J. Howie, *On locally indicable groups*, Math. Z. **180** (1982), no. 4, 445–461.
- [HS83] J. Howie and H. R. Schneebeli, *Homological and topological properties of locally indicable groups*, Manuscripta Math. **44** (1983), no. 3, 71 – 93.
- [Ju06] A. Juhász, *Holomorphic discs and sutured manifolds*, Algebraic & Geometric Topology **6** (2006), 1429–1457.
- [Ju08] A. Juhász, *Floer homology and surface decompositions*, Geometry and Topology **12:1** (2008), 299–350.
- [Ju08b] A. Juhász, *The sutured Floer homology polytope*, math.GT/0802.3415.
- [Ka05] O. Kakimizu, *Classification of the incompressible spanning surfaces for prime knots of 10 or less crossings*, Hiroshima Math. J. **35** (2005), no. 1, 47–92.
- [Ka96] A. Kawauchi, *A survey of knot theory*, Birkhäuser Verlag, Basel, 1996.
- [Mc02] C. T. McMullen, *The Alexander polynomial of a 3-manifold and the Thurston norm on cohomology*, Ann. Sci. Ecole Norm. Sup. (4) **35** (2002), no. 2, 153–171.
- [Ni24] J. Nielsen, *Die Isomorphismengruppe der freien Gruppen*, Math. Ann. **91** (1924), 169–209.

- [OS03] P. Ozsváth and Z. Szabó, *Heegaard Floer homology and alternating knots*, *Geom. Topol.* **7** (2003), 225–254.
- [OS04a] P. Ozsváth and Z. Szabó, *Holomorphic disks and topological invariants for closed 3-manifolds*, *Annals of Mathematics* **159** (2004), no. 3, 1027–1158.
- [OS04b] P. Ozsváth and Z. Szabó, *Holomorphic disks and 3-manifold invariants: properties and applications*, *Annals of Mathematics* **159** (2004) no. 3, 1159–1245.
- [OS04c] P. Ozsváth and Z. Szabó, *Knot Floer homology, genus bounds and mutation*, *Topology Appl.* **141** (2004), 59–85.
- [OS08a] P. Ozsváth and Z. Szabó, *Holomorphic disks, link invariants, and the multivariable Alexander polynomial*, *Algebr. Geom. Topol.* **8** (2008), no. 2, 615–692.
- [OS08b] P. Ozsváth and Z. Szabó, *Link Floer homology and the Thurston norm*, *J. Amer. Math. Soc.* **21** (2008), no. 3, 671–709.
- [Ra03] J. Rasmussen, *Floer homology and knot complements*, Harvard University thesis (2003), math.GT/0306378.
- [Pe08] T. Perutz, *Hamiltonian handleslides for Heegaard Floer homology*, math.GT/0801.0564
- [Sc89] M. Scharlemann, *Sutured manifolds and generalized Thurston norms*, *J. Differential Geom.* **29** (1989), no. 3, 557–614.
- [Th86] W. P. Thurston, *A norm for the homology of 3-manifolds*, *Mem. Amer. Math. Soc.* **59** (1986), no. 339, i–vi and 99–130.
- [Tu90] V. Turaev, *Euler structures, nonsingular vector fields, and Reidemeister-type torsions*, *Math. USSR-Izv.* **34** (1990), no. 3, 627–662.
- [Tu97] V. Turaev, *Torsion invariants of Spin^c -structures on 3-manifolds*, *Math. Res. Lett.* **4** (1997), no. 5, 679–695.
- [Tu98] V. Turaev, *A combinatorial formulation for the Seiberg-Witten invariants of 3-manifolds*, *Math. Res. Lett.* **5** (1998), no. 5, 583–598.
- [Tu01] V. Turaev, *Introduction to combinatorial torsions*, Birkhäuser Verlag, Basel, 2001.
- [Tu02] V. Turaev, *Torsions of 3-dimensional manifolds*, Progress in Mathematics, 208. Birkhäuser Verlag, Basel, 2002.
- [Wa78] F. Waldhausen, *Algebraic K-theory of generalized free products. I, II.*, *Ann. of Math.* (2) **108** (1978), no. 1, 135–204.

UNIVERSITY OF WARWICK, COVENTRY, UK
E-mail address: s.k.friedl@warwick.ac.uk

UNIVERSITY OF CAMBRIDGE, CAMBRIDGE, UK
E-mail address: A.Juhasz@dpms.cam.ac.uk

UNIVERSITY OF CAMBRIDGE, CAMBRIDGE, UK
E-mail address: J.Rasmussen@dpms.cam.ac.uk